

# MULTI-PHASE, MULTI-COMPARTMENT, CAPSULAR DELIVERY APPARATUS AND METHODS FOR USING SAME

## BACKGROUND

### 1. Related Applications

This Application is a continuation-in-part of PCT/US/03/10816 filed April 09, 2003, and entitled "MULTI-PHASE, MULTI-COMPARTMENT CAPSULAR SYSTEM", the contents of which are hereby incorporated by reference in their entirety. This application incorporates herein by reference U.S. Provisional Application Serial No. 60/371,448, filed April 10, 2002, and entitled "INTEGRATED CAPSULE DELIVERY APPARATUS AND METHOD." This application further claims the benefit of U.S. Application Serial No. 10/369,427, filed February 18, 2003, entitled "MULTI-PHASE, MULTI-COMPARTMENT CAPSULAR DELIVERY APPARATUS AND METHODS FOR USING SAME," which is hereby incorporated herein by reference. This application further claims the benefit of U.S. Application Serial No. 10/368,951, filed February 18, 2003, entitled "PROCESS FOR ENCAPSULATING MULTI-PHASE, MULTI-COMPARTMENT CAPSULES," which is hereby incorporated herein by reference. This application further claims the benefit of U.S. Application Serial No. 10/369,244, filed on February 18, 2003, and entitled "MULTI-PHASE, MULTI-COMPARTMENT CAPSULAR DELIVERY APPARATUS FOR THERAPEUTIC COMPOSITIONS AND METHODS FOR USING SAME," which is hereby incorporated herein by reference. This application further claims the benefit of U.S. Application Serial No. 10/369,247, filed February 18, 2003, and entitled "PROCESS FOR ENCAPSULATING MULTI-PHASE, MULTI-COMPARTMENT CAPSULES FOR THERAPEUTIC COMPOSITIONS," which is hereby incorporated herein by reference.

### 2. The Field of the Invention

The present invention relates to delivery of active ingredients or medicaments and, more particularly, to novel capsular delivery apparatus and methods for delivering one or

more active ingredients or medicaments having diverse physical states (*e.g.*, solid, liquid, gas or dispersion) into a single dosage, multi-compartment capsule.

The present invention further relates to methods for the administration of a plurality of heterogenous chemical and biological compounds to animals and humans using a multicompartment delivery system for treatment of different conditions or the same condition or diseases (different or same) in one or more organ systems.

### 3. Background of the Invention

As appreciated by those skilled in the art, the contemplation, design, testing and manufacture of chemicals and biomolecules for administration to humans and animals, as nutritional or therapeutic agents, requires a thorough integration of clinically contemplated delivery principles and modalities. Chemicals and biomolecules that may be administered to humans and animals are often referred to herein as "actives," "active ingredients" or "medicaments."

Oral administration has become one of the most frequent routes for delivering one or more active ingredients or medicaments to the body. Active ingredients or medicaments, such as nutritional or therapeutic agents, may be orally administered in a variety of physical states (*i.e.*, solid, liquid or gas). Tablets and capsules are generally the most common vehicle for the oral delivery of medicaments. As appreciated, a tablet may be broadly characterized as a compressed powder or granular solid. Prior to compression of the granular powder comprising the medicament into tablet form, the presence of one or more excipients may be required. An excipient includes any inert substance (*i.e.*, gum arabic, starch or the like) combined with a principal ingredient to facilitate the preparation of an agreeable or convenient dosage form of the active or medicament. Functional characteristics of excipients may include, for example, disintegration, lubrication, appearance, palatability, shelf-stability or the like.

Those skilled in the art also developed capsules as a contrivance for containing a solid or liquid dosage form of a medicament. Traditional capsular embodiments include a first containment section referred to as a base, and a second containment section referred to as a cap. The two pieces of the capsule are usually formulated and designed in a manner such that the material to be encapsulated may be introduced into the base section, whereas the open end of the cap section may be correspondingly positioned over the open end of the base. The walls of the cap and base are generally in physical contact with one another to form a single internal cavity. A means for structurally sealing the cap in relation to the base may also be incorporated during manufacture to insure non-tampering of the capsule. In this regard, those skilled in the art developed sealing technology which contemplates banding, heat fusion (spot-welding) and snap seals which utilize a "tongue and groove" scheme.

The outer walls of a capsule are preferably formed of a soluble ingredient, such as, for example, gelatin (animal-based product), starch, hydrophillic polymer or hydroxypropyl methyl-cellulose (HPMC), which provides a barrier for containing the active ingredient or medicament, in powder or liquid form, within the internal periphery of the capsule walls. Traditionally, hard gelatin capsules may be manufactured by dipping plates of stainless steel pins into a pool of gelatin solution. The pins are then removed from the gelatin and rotated while the gelatin is dried in a kiln with forced, humidity-controlled air. Once dried, the gelatin capsules are typically stripped from the pins, trimmed to a suitable length and then joined together (*e.g.*, base and cap) and packaged for production use.

With the advent of automated encapsulation machinery, the responsibility to produce encapsulated products shifted mainly to industrial manufacturers. Contemporaneous with the development of the encapsulation industry, those skilled in the art have advanced the state of the encapsulation art. For example, several significant improvements in encapsulation technology have been seen over the last forty years. These technological improvements have included, for example, the development of soft elastic capsules, film-coating techniques, micro-encapsulation and multiple-compartment technology.

Soft elastic capsules, often referred to as soft gelatin capsules, were developed in an effort to provide means for encapsulating liquids and other medicaments which are typically poorly soluble in water. In preferred design, soft elastic capsules are made from a thicker and more plastic gelatin having an increased flexibility due to the addition of a polyol, such as glycerin or sorbitol. The addition of such plasticizers has been found, however, to have the potential disadvantage of increasing the risk for microbial growth. Thus an antimicrobial, such as a paraben or sorbic acid, may be added to the soft elastic capsule shell in order to address any microbial concern.

Prior art film-coating techniques generally involve a plating process, whereby a thin, uniform film may be deposited onto the outer surface of the delivery vehicle (*e.g.*, tablet or capsule). Several successive layers may be deposited onto the outer surface of the vehicle, if desired, in an effort to facilitate various desirable properties. For example, sugar-coating, a precursor to film-coating, has been used by those skilled in the art for more than one hundred years to make tablets more palatable. Other advantages or properties of film-coating may include for example, but not by way of limitation, protection from moisture, oxidation, controlling microbial contamination and inhibiting modification of the chemical properties of the active ingredient. As further appreciated by those skilled in the art, prior art film-coating may form an interfacial barrier between two chemicals or chemical compounds that might otherwise react when they come into contact.

Enteric coatings and sustained-release formulations are contemplated as variations on prior art film-coating techniques. In particular, enteric coating describes a process where the delivery vehicle (*e.g.*, tablet or capsule) is coated with one or more layers of chemicals that are somewhat resistant to extreme pH conditions. For example, conditions of extremely low pH are commonly encountered in the stomach. Many active ingredients or medicaments are in the form of a pharmaceutical salt and thus highly susceptible to ionization in the presence of hydrogen ions. Thus, the presence of an enteric coating generally provides a level of protection as to degradation of the active ingredient or medicament until transit from the stomach into the small intestine is accomplished.

Film coatings have also led to the development of delivery vehicles (*e.g.*, tablets and capsules) having sustained-release properties. Mixtures of waxes, cellulose, silicone and similar resins have been found useful by those skilled in the art for creating-sustained release coatings. In principle, these prior art coatings function to delay the release of the active ingredient or medicament to the targeted body system, thereby facilitating a timed, absorption rate in the body. Furthermore, the entire daily dosage of an active or medicament may be contained in a single, sustained-release delivery vehicle (*e.g.*, tablet or capsule), whereas the immediate absorption of the entire dosage could possibly lead to an overdosage of the medicament. Thus, by layering quanta of medicament with differential coatings, the dosage undergoes a controlled release over specified time period. The application of sustained-release film coating technology therefore may inherently facilitate the delivery of a total daily dosage amount of an active or medicament to be released to the body in controlled increments.

Over the last several years, a considerable amount of attention has been focused on the further development of multi-compartment capsule technology for the delivery of therapeutic and diagnostic agents. Series formulations teach the use of membranes or other types of barriers to cordon a line of separate chambers within a single encapsulating shell. As appreciated, the purpose of such multi-compartment delivery devices is the administration of multiple dosages. Moreover, multiple-compartment delivery mechanisms of the prior art were developed to circumvent or diminish the effects of harsh pH environments within humans. For example, the prior art contemplates a hard capsule formulation which contains three different compartments of active medicaments for administration to the vaginal and rectal areas. In preferred structure, the formulation outer, rapid-release layer may contain an active medicament and excipient; the middle, intermediate-release layer may include a powder form of active medicament; and the inner, slow-release layer may contain pellets or granules of active medicament.

Also taught in the prior art are multi-compartment capsules having groups of spheroids with pH-dependent coatings which are encapsulated within a hard gelatin shell and

provided for treating female yeast infection. The first spheroid is preferably uncoated and may be in a powder form; the second spheroid may contain a pH sensitive coat; and the inner spheroid may include a pH insensitive coat.

5 In addition to pH-sensitive coatings, hydrogels and other gastric retention technologies have been developed by those skilled in the art in an effort to retard the progression of the delivery vehicle during enteric transit. This retarding action, presumably, allows the full amount of active medicament to be released and/or targeted to a specific area of the gastrointestinal tract. Hydrogel and related gastric retention devices of the prior art generally rely upon the imbibing of water into a center core which is filled with cellulose or  
10 similar water absorbent material. In preferred operation, the material swells and releases multiple compartments of active medicament. The concept of using bulk size to slow transit of single active medicament in a single physical state is thus appreciated.

In an effort to administer active ingredients or medicaments to a specific location in the body to treat a specific disorder caused by a specific pathogen, those skilled in the art  
15 have used targeted-release systems using multi-compartment capsular technology. For example, a method for carrying out a triple therapy against the microorganisms *Helicobacter pylori*, a known infectious agent which is believed largely responsible for the development of gastric ulcer disease, was developed which comprises the steps of oral administration of a pharmaceutical dosage form comprising an internal capsule placed inside an external  
20 capsule, wherein the external capsule comprises a soluble salt of bismuth and a first antibiotic, and the internal capsule comprises a second antibiotic. In addition, multi-compartmental capsules were developed which combine a nutrient supplement with a viable direct-fed microbial (*i.e.*, gastrointestinal microorganisms, including bacteria, live cell yeasts, fungi or a combination thereof) for the purpose of treating livestock for feeding  
25 disorders and improving feed efficiency.

A disadvantage with prior art encapsulation technology is when the base and corresponding cap of a capsule are joined, dead space volume is typically created within the internal periphery of the capsule. Internal capsular dead space may be filled with an air

bubble which may ultimately react with one or more of the active ingredients or medicaments introduced within the capsule, thereby potentially degrading the quality and effectiveness of the active ingredients.

Although the prior art discloses multiple compartment, capsular delivery technology, these manifestations generally includes one of two approaches. For example, one approach contemplates the introduction of a single active or medicament into multiple capsular compartments to vary the temporal release of the medicament and ultimately the absorption rate into the body. Another approach contemplates the introduction of a plurality of active ingredients or medicaments into different compartments of a single capsule for delivery to a specific area of the body to treat a targeted illness or condition.

The use or contemplation of multiple-compartment capsular delivery apparatus or methods which deliver different physical forms of the same active or medicament, or a variation in physical forms of different actives or medicaments in a single dosage, however, has not heretofore been contemplated in the art. As appreciated by those skilled in the art, active ingredients or medicaments may take the physical form of a solid (*e.g.*, pill, tablet, capsule (both hard and soft elastic), powder, granulation, flakes, troches (lozenges and pastilles), suppositories and semi-solid ointments, pastes, emulsions and creams), a liquid (*e.g.*, solution, spirits, elixir, syrups, sprays and fluid extracts), a gas or a dispersion. A dispersion is a system in which a dispersed phase is distributed through a continuous phase (*e.g.*, aerosols (liquid or solid in gas), suspensions (solid in liquid), emulsion (liquid in liquid), foam (gas in liquid), solid foam (solid in gas) or gel (liquid or solid in solid)). Dispersions can be classified as molecular, colloidal and coarse, depending on size. In many circumstances, however, the different physical forms or phases of more than one active ingredient or medicament may not be suitably combined or mixed together without altering the individual desirable properties of the active ingredient or medicament. For example, although it would be possible and desirable to formulate a dispersion by combining a first active ingredient in the solid state with a second active ingredient that exists as a liquid, adverse chemical interactions between the active ingredients may adversely affect various characteristics of the ingredients, including but not limited to, their shelf lives. The resulting chemical decomposition – and the potential formation of any unwanted side products – could result in diminished drug potency or even toxicity to a patient.

Additionally, the physical properties of crystalline active ingredients could be drastically altered in scenarios where it is desirable to co-administer a crystalline active ingredient with a liquid or semi-liquid different active ingredient. In this context, the control of physical properties such as active ingredient dissolution rate and solubility is often a critical factor in determining the overall bioavailability of the active ingredient. It is well established in the art that different polymorphs or solvates of the same crystalline active ingredient exhibit dramatically different solubility and dissolution rates. Thus, combining a crystalline active agent with a liquid or semi-liquid active agent could give rise to an equilibrium between concentrations of different polymorphs and/or solvates of the crystalline active ingredient, and thereby frustrate efforts at tailoring an active ingredient mixture to its intended purpose as a medicament.

Another shortcoming with co-administering plural active ingredients in different physical forms in an intimate mixture is the potential for adverse *in vivo* drug-drug interactions upon administration. The desire to co-administer these active ingredients would be offset by the one active ingredient, for example as in a liquid or semi-liquid (e.g., a paste, solution, or syrup) form, becoming rapidly available. In this context, the active ingredient may adversely react with a co-administered drug, for example a less bioavailable solid or semi-solid, in a physiological environment. Thus, the true therapeutic benefit resulting from the pharmacological effects of the individual active agents may never be realized. It would be desirable to co-administer plural active ingredients while insuring against the potential of such harmful drug-drug interactions.

Providing active ingredients or medicaments in separate capsules may also be undesirable in the context of patient compliance. Geriatric and pediatric populations in particular disfavor the handling and consumption of multiple capsules of active ingredients. Patient compliance is essential in maintaining patient health in many dosage regimens. For example, deviations from accurate dosing and consistent consumption of immunosuppressant therapies can result in severe or even lethal consequences for a patient. Providing combined dosages of active ingredients would result in fewer capsules a patient or consumer would have to take, and thereby contribute to an overall increase in compliance.

Therefore, it would be desirable to provide a multi-compartment capsular delivery apparatus and methods that provide active ingredients or medicaments having diverse physical properties (e.g., solid, liquid, gas or dispersion), which may or may not be properly



combined or stored together into a unitary structure (i.e., multi-compartment capsule) for usage in a single dosage form. The present invention, in overcoming the shortcomings of the prior art, satisfies these and other objectives.

\* \* \* \* \*

5           The art and practice of pharmacy can be divided into four distinct divisions. Pharmacology is the study of interactions occurring between the pharmacologic agent, or medicament and specific targeted cells in the body. More specifically, the interaction between an active agent and a cellular receptor along with the resulting change in cell physiology is examined. Medicinal chemistry is largely concerned with the identification of  
10   naturally occurring and synthetic compounds which possess medicinal characteristics.

Pharmacotherapeutics is the holistic application of pharmacy practice to specific pathologies, illnesses, and other body functions. Finally, Pharmaceutical science ascertains or regulates the composition of medicinal substances, and is largely directed to the development of new mechanisms for delivering chemicals and biomolecules into animals and  
15   humans. A subcategory of pharmaceutical science is called pharmacokinetics and sometimes generally referred to as biopharmaceutics.

A.D.M.E. is an acronym often used to describe the four essential components to pharmaceutical science: absorption, distribution, metabolism, and elimination, respectively. One way to differentiate between pharmacology and pharmaceutical science is that the  
20   former is primarily concerned with the effect of the medicament on the body, whereas, the latter is primarily concerned with the delivery and time-course of the medicament on its journey through the body.

In clinical applications, chemicals and biomolecules are often referred to as active ingredients or medicaments. Medicaments may include “pharmaceuticals, nutraceuticals,  
25   biotechnicals, vitamins, minerals and dietary supplements.” Oral administration is the most frequent route for delivery of medicaments. Medicaments may be orally administered in a variety of physical states, including, solid, liquid, dispersion, and gaseous forms. As appreciated, tablets and capsules are the most common vehicle for oral delivery of medicaments.

Frequently, a medical or surgical patient may receive a plurality of concurrent medicaments. Data has been accumulated to demonstrate that patients undergoing a surgical procedure may receive ten (10) or more medicaments during the surgery and the resulting surgical recovery period. Some patients who have undergone organ transplantation or who have contracted human immunodeficiency virus (HIV) may receive three (3) or more medicaments which require multiple administrations per day. HIV patients often receive many more than three (3) medicaments. These medicaments may be necessary for the treatment of several conditions occurring in a plurality of organ systems or they may be necessary to treat a single condition or some combination thereof.

In some cases, it may be desirable to combine a plurality of medicaments because of a synergistic interaction between a plurality of medicaments. This synergy may enhance the efficacy of one or more of the medicaments. Medicaments may be combined to increase the intensity of response or efficacy. A plurality of medicaments, in combination, may be homergic (*i.e.*, elicit the same quality of effect). In many cases, a plurality of homergic medicaments may also be homodynamic (*i.e.*, interact with the same receptor). A plurality of homergic medicaments may be additive, supra-additive and infra-additive. A plurality of combined medicaments which do not produce the same quality of response may be called, heterergic. When heterergy is found to be a positive effect (*i.e.*, at least one medicament enhances the response to another medicament), this may be called synergism and is sometimes called synergy.

In further cases, it may be desirable to combine a plurality of medicaments to decrease their individual dosages and possibility for toxicity. It may also be desirable to combine a plurality of medicaments to target the treatment of a disease, illness or condition from divergent angles. It may be desirable to combine a plurality of medicaments to minimize the side effects and adverse effects of one or more medicaments. It may be still further desirable to combine a plurality of medicaments to alter the pharmacokinetic characteristics of one or more medicaments. For example, alterations in the absorption, distribution, metabolism or elimination of one or more medicaments.

Fixed combinations of a plurality of medicaments have been generally disfavored due to any number of perceived disadvantages. These disadvantages may include, for example: (1) complicating the interpretation of safety and efficacy in therapeutic regimens, (2) there may be inter-patient differences to fixed combinations, (3) there may be difficulties in dosage

titration, and (4) the delivery platforms for fixed combinations have generally been found to be uneconomical to produce.

On the other hand, fixed combinations of a plurality of medicaments may lead to several therapeutic advantages, including, for example, but not by way of limitation: (1) increasing patient compliance with therapy, (2) increasing efficacy by optimizing timing of medicaments, (3) minimization of side effects and adverse effects, (4) enhancement of pharmacokinetic characteristics of one or more medicaments in a fixed combination, (5) increased patient quality of life, (6) optimization of institutional resources by minimizing the amount of medicament administrations, and (7) minimizing patient length of stay in institutional facilities by optimizing therapy.

Prior art therapeutic technologies contain isolated examples of pharmaceutical formulations containing fixed combinations of medicaments. However, therapeutic technologies of the prior art teach a fixed combination, wherein a plurality of medicaments are placed into a single receiving chamber in the delivery formulation (i.e., no separation between the plurality of medicaments).

In view of the state of the technology as it exists today, generally, therapeutic apparatus and methods are needed to provide a plurality of medicaments for medical and surgical conditions, as well as maintenance of normal health function for delivery to animals and humans using a multi-chambered delivery apparatus. Such apparatus and methods for delivering a plurality of medicaments to animals and humans using a multi-chambered delivery apparatus are contemplated herein.

#### BRIEF SUMMARY AND OBJECTS OF THE INVENTION

In view of the foregoing, it is a primary object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering diverse physical states (*e.g.*, solid, liquid, gas or dispersion) of a single active ingredient or medicament, or a plurality of active ingredients or medicaments, in a single dosage form, wherein at least two of the active ingredients or medicaments have physical states that differ.

It is also an object of the present invention to provide novel integrated capsule delivery apparatus and methods which facilitate various desirable properties including, for example, controlling time-release of key active ingredients or medicaments, prolonging shelf-life of the active ingredients or medicaments, improving palatability, reducing overall production costs and, accordingly, reducing the number of capsules consumed by a patient or consumer as nutritional or therapeutic agents.

Further, it is an object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule having one or more active ingredients in a primary capsule, and one or more active ingredients introduced into a secondary smaller capsule having a size sufficient for being selectively positionable within the primary capsule, wherein the active ingredient(s) within the primary capsule comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in the secondary capsule.

It is an additional object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule having one or more active ingredients in a primary capsule and the same active ingredient(s) introduced into a smaller secondary capsule having a size sufficient for being positionable within the primary capsule, wherein the active ingredient(s) in the primary capsule comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the active ingredient(s) in the secondary capsule.

It is a further object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule wherein at

least one of the primary and secondary capsules include a time-release coating for controlling the release of the active ingredient(s) contained therein.

It is also another object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (e.g., pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule having one or more active ingredients in the capsular body, wherein the capsule includes a longitudinally extending body and at least one dividing wall formed along a length of the extending body to form a first chamber and an opposing second chamber within the capsular body and introducing at least one active ingredient or medicament having a first physical state into the first chamber and at least one active ingredient or medicament having a second physical state into a second chamber, whereas the physical state (e.g., solid, liquid, gas or dispersion) of the ingredient(s) in the first chamber is different from the physical state of the ingredient(s) in the second chamber.

It is an additional object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (e.g., pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule having a longitudinally extending body and one or more dividing walls disposed along the length of the longitudinally extending body of the capsule, wherein the capsule and one or more of the dividing walls contained therein may include time-release coatings for controlling the release of the active ingredients or medicaments contained therein, respectively.

It is a further object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (e.g., pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule having a plurality of active ingredients or medicaments having the physical form of a solid (e.g., pill, tablet, capsule (both hard and soft elastic), powder, granulation, flakes, troches (lozenges and

pastilles), suppositories and semi-solid ointments, pastes, emulsions and creams), a liquid (*e.g.*, solution, spirits, elixir and fluid extracts), a gas or a dispersion (*e.g.*, aerosols (liquid or solid in gas), suspensions (solid in liquid), emulsion (liquid in liquid), foam (gas in liquid), solid foam (solid in gas) or gel (liquid or solid in solid), wherein the physical form of the active ingredients differ between a primary and secondary capsule, and between one or more dividing walls disposed in spaced-apart relationship along the length of a longitudinally extending capsular body.

It is a still further object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) in the form of a single dosage, multi-compartment capsule, wherein an encapsulation process comprises the steps of: (1) providing a capsule comprising a first end, a second end, a longitudinally extending body having a length disposed between the first and second ends, and a plurality of dividing walls spaced apart along the length of the extending body, wherein the dividing walls form a plurality of receiving chambers; (2) introducing at least one active ingredient having a first physical state into a first receiving chamber; (3) introducing at least one active ingredient having a second physical state into a second receiving chamber; (4) introducing at least one active ingredient having a third physical state into a third receiving chamber, wherein the physical states of at least two of the active ingredients introduced into the first, second or third receiving chambers differ; and (5) sealing the first and second ends of said capsule.

Additionally, it is an object of the present invention to provide novel integrated capsule delivery apparatus and methods for delivering a single dosage, multi-compartment capsule comprising a capsular base and cap configuration, wherein the size and shape of the cap, relative to its sealing relationship with the base, generally eliminates or substantially reduces any potential dead space volume within the internal periphery of the capsule, thereby functionally negating the opportunity for reaction between an air bubble and one or more

active ingredients introduced into the capsule and, accordingly, improving stability of the capsular ingredient(s).

Consistent with the foregoing objects, and in accordance with the invention as embodied and broadly described herein, one presently preferred embodiment of the novel integrated capsule delivery apparatus and methods of the present invention comprises a multi-compartment capsule including a primary capsule and a secondary capsule selectively positionable within an internal periphery of the primary capsule. The secondary capsule may include a base, a corresponding cap and one or more receiving chambers. Each of the receiving chambers of the secondary capsule may be formed having an internal periphery sufficient for receiving at least one active ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. Similarly, the primary capsule may be formed having a base, a corresponding cap and one or more receiving chambers. The receiving chambers of the primary capsule may be formed having an internal periphery sufficient for receiving the secondary capsule and one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a physical state (*i.e.*, solid, liquid, gas or dispersion) different from the physical state of the active ingredient(s) housed within the receiving chamber of the secondary capsule.

As further contemplated herein, a multi-compartment capsule is provided comprising a base, a corresponding cap and one or more dividing walls positionable between the base and the cap. Structurally, the size, shape and positioning of the dividing walls relative to the base and corresponding cap facilitates the formation of at least two, independent and separate receiving chambers. Each of the receiving chambers having an internal periphery sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. In preferred design, the physical state (*e.g.*, solid, liquid, gas or dispersion) of the active ingredient(s) in the first receiving chamber is different from the physical state of the active ingredient(s) in the second receiving chamber. After introducing one or more active

ingredients or medicaments into each receiving chamber, the cap may be selectively positioned in sealing relationship with the base to form one presently preferred embodiment of the single, dosage multi-compartment capsule.

One presently preferred embodiment of an encapsulation process for forming a multi-compartment capsule may comprise the steps of: (1) providing a primary capsule having a base, a corresponding cap and a receiving chamber; (2) providing a secondary capsule having a base, a corresponding cap and a receiving chamber; (3) introducing at least one ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a first physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber of the secondary capsule and selectively positioning the cap in sealing relationship with the base; (4) introducing at least one ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a second physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber of the primary capsule, wherein the first physical state of the ingredient(s) in the secondary capsule is different from the second physical state of the ingredient(s) in the primary capsule; and (5) introducing the secondary capsule into at least a portion of the receiving chamber of the primary capsule and selectively positioning the cap in sealing relationship with the base to form a single dosage multi-compartment capsule.

In alternate presently preferred embodiments of the present invention, a tertiary capsule comprising a base, a corresponding cap and a receiving chamber having an internal periphery sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) may be selectively introduced within an internal periphery of at least one receiving chamber of the secondary capsule. After the introduction of at least one active ingredient into one or more receiving chambers of a tertiary capsule pursuant to an encapsulation process of the present invention, the cap of the tertiary capsule may be selectively positioned in sealing relationship with the base and then introduced into at least a



portion of the internal periphery of the secondary capsule, together with one or more active ingredients therein. It is contemplated herein that at least two of the active ingredients introduced within the receiving chambers of the primary, secondary and tertiary capsules, respectively, comprise at least two different physical states (e.g., solid, liquid, gas or dispersion).

In preferred structural design, the primary capsule may comprise a cap having a generally U-shaped configuration adapted to provide a sealing relationship when engaging the corresponding base, thereby reducing dead space volume in the internal periphery of the cap and receiving chamber of the base. A cap having a configuration adapted to generally eliminate or substantially reduce potential dead space volume of the cap and receiving chamber of the base may, accordingly, function to negate the potential for a reaction between an air bubble and one or more active ingredient(s) introduced into the base of the primary capsule.

Alternatively, a multi-compartment capsule of the present invention may include the introduction of a filling material into the cap of the primary capsule, the cap having a general cylindrical configuration adapted to provide a sealing relationship when engaging the corresponding base. An amount of filling material may be introduced into at least a portion of the internal periphery of the cap to fill, either partially or completely, the inner volume of the cap, thereby reducing the dead space volume in the cap and the internal periphery of the receiving chamber of the base. In this regard, the introduction of a filling material relative to the internal periphery of the cap may generally eliminate or substantially reduce the potential dead space volume, thus functionally negating the potential for a reaction between an air bubble and one or more active ingredient(s) introduced into the base of the primary capsule.

The primary, secondary or tertiary capsules, in accordance with the present invention, may be formed having the same or different colors. Moreover, the base and corresponding cap of a single capsule may be formed having different colors in an effort to enhance the aesthetics of the capsule to the consumer. In one presently preferred embodiment of a multi-compartment capsule of the present invention, the dosage may be banded, sealed or

easily dividable in a contact area of the primary and secondary capsules or the sealing band may be color-coded to assist in branding, if desired.

It is further contemplated herein that a multi-compartment capsule of the present invention may comprise component parts of the capsule having various time-release coatings to facilitate the release and ultimately the absorption of those active ingredients introduced into the different receiving chambers of the multi-compartment capsule to release at different release rates. In particular, a primary capsule may be formed having a conventional time-release coating that dissolves and releases the active ingredient(s) contained therein before the timed-release of the active ingredient(s) contained within a secondary capsule.

Likewise, the dividing walls disposed within the internal periphery of the base of a capsule may be formed having conventional time-release coatings that dissolve and release the active ingredients within each receiving chamber defined by the dividing walls at different rates, thereby delivering the active ingredients or medicaments contained within a multi-compartment capsule at different rates. Certain active ingredients or medicaments may, therefore, be delivered at a selected interval, while other ingredients may be released at a later interval. In this way, the novel design of the multi-compartment capsules of the present invention may facilitate precision delivery of active ingredients to targeted areas of the consumer.

\* \* \* \* \*

Still further, a primary object of the present invention is to provide novel delivery apparatus and methods for affecting multiple organ systems in animals or humans using a plurality of medicaments delivered by a pharmaceutical formulation comprising a multi-chambered apparatus. Accordingly, the present invention provides novel delivery apparatus and administration techniques or methods aimed at affecting multiple organ systems in an animal or human using a plurality of medicaments. A delivery apparatus may be in any multi-chambered apparatus, but preferably in a capsular formulation. Thus, a plurality of medicaments may be encapsulated and stored separately within a larger capsule until the time of ingestion, consumption, or the like. Upon consumption, the capsule walls of one or more dividing walls of a capsule may dissolve to release their contents. Different methods of

encapsulation may be used to deliver their respective contents, including but not limited to, dissolution, melting, ablation or biodegradation of the encapsulating wall. In certain embodiments and as contemplated herein, the medicaments retained in the multicompartment capsule may actually diffuse through one or more of the encapsulating walls.

5

\* \* \* \* \*

In one embodiment of the present invention there is a multi-compartment capsule, comprising a first receiving chamber comprising at least one ingredient having a first physical state, wherein said ingredient is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; and a second receiving chamber  
10 comprising at least one ingredient having a second physical state, wherein said ingredient is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; said first physical state of said ingredient of said first receiving chamber being different from said second physical state of said ingredient of said second receiving chamber.

In another embodiment of the present invention, there is a multi-compartment capsule  
15 as defined above, further comprising a base and a corresponding cap, wherein said cap is configured to provide a sealing relationship when engaging said base.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said cap comprises a configuration adapted to reduce dead volume space within said first receiving chamber.

20 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, further comprising a filling material introduced into said cap to reduce dead volume space within said first receiving chamber.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said filling material is selected from the group consisting of  
25 gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polyvinylacetate-phtalate, polymerisates of acrylic or methacrylic esters and combinations thereof.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said first receiving chamber comprises no dead volume space.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said first receiving  
5 chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said second receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule  
10 as defined above, wherein said solid is selected from the group consisting of a pill, a tablet, a capsule, a powder, granulation, flakes, a troche, a suppository, an ointment, a paste, an emulsion and a cream.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said liquid is selected from the group consisting of a solution, a  
15 spirit, an elixir, a spray, a syrup and a fluid extract.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said dispersion is selected from the group consisting of an aerosol, a suspension, an emulsion, a foam, a solid foam and a gel.

In another embodiment of the present invention, there is a multi-compartment capsule  
20 as defined above, wherein said first receiving chamber comprises a time-release coating.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said second receiving chamber comprises a time-release coating.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said time-release coating of said second receiving chamber is  
25 different from said time-release coating of said primary capsule.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, further comprising a third receiving chamber comprising at least one ingredient.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said third receiving chamber is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said third receiving chamber comprises a physical state selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said third receiving chamber comprises a time-release coating.

10 In another embodiment of the present invention, there is a multi-compartment capsule, comprising a primary capsule comprising at least one ingredient having a first physical state, wherein said ingredient is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; a secondary capsule comprising at least one ingredient having a second physical state, wherein said ingredient is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral;  
15 said first physical state of said ingredient of said primary capsule being different from said second physical state of said ingredient of said secondary capsule; and said primary capsule comprising an internal periphery sufficient for receiving said ingredient and said secondary capsule therein.

20 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule further comprises a base and a corresponding cap, wherein said cap is configured to provide a sealing relationship when engaging said base.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule comprises no dead volume space.

25 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said first physical state of said ingredient in said primary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said second physical state of said ingredient in said secondary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule comprises a time-release coating.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said secondary capsule comprises a time-release coating.

10 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said time-release coating of said secondary capsule is different from said time-release coating of said primary capsule.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said third receiving chamber comprises a time-release coating.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, oleoresin, polyvinylacetate, hydroxypropyl methyl cellulose, polymerisates of acrylic or methacrylic esters, polyvinylacetate-phtalate and combinations  
20 thereof.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

25 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said soft elastic capsule includes an antimicrobial selected from the group consisting of paraben and sorbic acid.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said secondary capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein,

alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, oleoresin, polyvinylacetate, hydroxypropyl methyl cellulose, polymerisates of acrylic or methacrylic esters, polyvinylacetate-phtalate and combinations thereof.

5           In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said primary capsule comprises a moisture content in the range of about 0% to 6% by weight.

          In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said secondary capsule comprises a  
10       moisture content in the range of about 0% to 6% by weight.

          In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary and secondary capsules contain at least one pharmaceutically acceptable lubricant in the range of about 0% to 10% by weight.

          In another embodiment of the present invention, there is a multi-compartment capsule  
15       as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and mixtures thereof.

          In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary and secondary capsules have different colors.

20           In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule is formed having a first color.

          In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said secondary capsule is formed having a second color different from said first color of said primary capsule.

25           In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule further comprises a base and a corresponding cap, wherein said cap is configured to provide a sealing relationship when engaging said base.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said base and said cap are formed having different colors.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said sealing relationship between said base and corresponding cap comprises no dead volume space.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said first receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

10 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said second receiving chamber capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule comprises a time-release coating.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said dividing wall comprises a time-release coating.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said time-release coating of said dividing wall is different from said time-release coating of said capsule.

20 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said third receiving chamber comprises a time-release coating.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, 25 gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, oleoresin, polyvinylacetate, hydroxypropyl methyl cellulose, polymerisates of acrylic or methacrylic esters, polyvinylacetate-phtalate and combinations thereof.



In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said dividing wall is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephthalate-acetate, oleoresin, polyvinylacetate, hydroxypropyl methyl cellulose, polymerisates of acrylic or methacrylic esters, polyvinylacetate-phtalate and combinations  
10 thereof.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said first receiving chamber comprises a moisture content in the range of about 0% to 6% by weight.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said second receiving chamber comprises a moisture content in the range of about 0% to 6% by weight.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule contains at least one pharmaceutically acceptable lubricant in the range of about 0% to 10% by weight.

20 In another embodiment of the present invention there is, an encapsulation process for forming a multi-compartment capsule, said process comprising the steps of providing a primary capsule having a base and a cap; providing a secondary capsule having a base and a cap; introducing at least one ingredient having a first physical state into said secondary capsule, wherein said ingredient introduced into said primary capsule is selected from the  
25 group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; positioning said cap of said secondary capsule in sealing relationship with said base; introducing at least one ingredient having a second physical state into said primary capsule, wherein said ingredient introduced into said primary capsule is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; and wherein said  
30 first physical state of said ingredient of said secondary capsule is different from said second

physical state of said ingredient of said primary capsule; introducing said secondary capsule into said base of said primary capsule; and positioning said cap of said primary capsule in sealing relationship with said base.

5 In another embodiment of the present invention there is, an encapsulation process as defined above, further comprising the step of reducing dead volume space within said primary capsule.

In another embodiment of the present invention, an encapsulation process as defined above, further comprising the step of introducing a filling material into said cap of said primary capsule to reduce dead volume space.

10 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said filling material is selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polyvinylacetate-phthalate,  
15 polymerisates of acrylic or methacrylic esters and combinations thereof.

In another embodiment of the present invention, an encapsulation process as defined above, wherein said cap of said primary capsule comprises a configuration sufficient for reducing dead volume space within the primary capsule.

20 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said primary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said secondary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

25 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient introduced into said primary capsule is the same as said ingredient introduced into said secondary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary capsule comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said secondary capsule comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said time-release coating of said secondary capsule is different from  
5 said time-release coating of said primary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the steps of providing a tertiary capsule having a base and a cap; introducing at least one ingredient having a third physical state into said tertiary capsule; positioning said cap of said secondary capsule in sealing relationship with said base;  
10 and introducing said tertiary capsule into said base of said secondary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said tertiary capsule is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as  
15 defined above, wherein said ingredient in said tertiary capsule comprises a physical state selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said tertiary capsule comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as  
20 defined above, wherein said primary capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polymerisates of acrylic or mthacrylic esters, polyvinylacetate-phtalate and combinations  
25 thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said secondary capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polymerisates of acrylic or mthacrylic esters, polyvinylacetate-phtalate and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said secondary capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient introduced in said primary capsule comprises a moisture content in the range of about 0% to 6% by weight.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said secondary capsule comprises a moisture content in the range of about 0% to 6% by weight.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary and secondary capsules contain at least one pharmaceutically acceptable lubricant in the range of about 0% to 10% by weight.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary and secondary capsules are formed having different colors.

In another embodiment of the present invention, there is an encapsulation process for forming a multi-compartment capsule, said process comprising the steps of providing a capsule comprising a cap, a base configured having a longitudinally extending body

including a length and at least one dividing wall formed along said length of said extending body, said dividing wall adapted to form a first receiving chamber and a second receiving chamber; introducing at least one ingredient having a first physical state into said second receiving chamber, wherein said ingredient introduced into said primary capsule is selected  
5 from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral; introducing at least one ingredient having a second physical state into said first receiving chamber, wherein said ingredient introduced into said primary capsule is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral, and wherein said first physical state of said ingredient of said second receiving chamber being  
10 different from said second physical state of said ingredient of said first receiving chamber; and positioning said cap in sealing relationship with said base.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of reducing dead volume space within said primary capsule.

15 In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of introducing a filling material into said cap to reduce said dead volume space.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said filling material is selected from the group consisting of gelatin,  
20 starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polyvinylacetate-phtalate, polymerisates of acrylic or methacrylic esters and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as  
25 defined above, wherein said cap comprises a configuration sufficient for reducing dead volume space within said capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said second receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

5 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said first receiving chamber comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said second receiving chamber comprises a time-release coating.

10 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said time-release coating of said second receiving chamber is different from said time-release coating of said first receiving chamber.

15 In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the steps of positioning a second dividing wall along said length of said extending body, said second dividing wall adapted to form a third receiving chamber; and introducing at least one ingredient having a third physical state into said third receiving chamber.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said third receiving chamber is selected from the group consisting of a nutraceutical, a vitamin, a dietary supplement and a mineral.

20 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said third receiving chamber comprises a physical state selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said dispersion is selected from the group consisting of an aerosol, a suspension, an emulsion, a foam, a solid foam and a gel.

25 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said third receiving chamber comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said capsule is formed of a material selected from the group

consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin, polymerisates of acrylic or methacrylic esters, polyvinylacetate-phthalate and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary and secondary capsules contain at least one pharmaceutically acceptable lubricant in the range of about 0% to 10% by weight.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said base and said cap of said capsule are formed having different colors.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of introducing two or more dividing walls adapted to form a plurality of receiving chambers into said base of said capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of introducing a capsule into one of said plurality of receiving chambers.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said capsule may comprise a multi-compartment capsule.

In another embodiment of the present invention, there is a multi-compartment capsule, comprising a first receiving chamber comprising at least one ingredient having a first physical state; and a second receiving chamber comprising at least one ingredient having a

second physical state, wherein said first physical state of said ingredient of said first receiving chamber being different from said second physical state of said ingredient of said second receiving chamber.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said first receiving chamber comprises no dead space.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said cap is configured to reduce dead volume space within said first receiving chamber.

10 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, further comprising a filling material introduced into said cap to reduce dead volume space within said first receiving chamber.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said first receiving chamber is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said second receiving chamber is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

20 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber comprises a pharmaceutical.

25 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber is selected from the group consisting of a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.



In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said first receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said physical state of said ingredient in said second receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said time-release coating of said second receiving chamber is different from said time-release coating of said primary capsule.

10 In another embodiment of the present invention, there is a multi-compartment capsule, comprising a primary capsule comprising at least one ingredient having a first physical state; a secondary capsule comprising at least one ingredient having a second physical state; said first physical state of said ingredient of said primary capsule being different from said second physical state of said ingredient of said secondary capsule; and  
15 said primary capsule comprising an internal periphery sufficient for receiving said ingredient and said secondary capsule therein.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule further comprises a base and a corresponding cap, wherein said cap is configured to provide a sealing relationship when engaging said  
20 base.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule comprises no dead volume space.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said primary capsule is selected from the group  
25 consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said secondary capsule is selected from the

group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber comprises a pharmaceutical.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said primary capsule comprises a moisture content in the range of about 0% to 6% by weight.

10 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient introduced in said secondary capsule comprises a moisture content in the range of about 0% to 6% by weight.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary and secondary capsules contain at least one pharmaceutically acceptable lubricant in the range of about 0% to 10% by weight.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and mixtures thereof.

20 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary and secondary capsules have different colors.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said primary capsule is formed having a first color.

25 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said secondary capsule is formed having a second color different from said first color of said primary capsule.

In another embodiment of the present invention, there is a multi-compartment capsule, comprising a capsule comprising a longitudinally extending body having a length; at

least one dividing wall formed along said length of said extending body, said dividing wall forming a first receiving chamber and a second receiving chamber; said first receiving chamber comprising at least one ingredient having a first physical state; said second receiving chamber comprising at least one ingredient having a second physical state; and said first  
5 physical state of said ingredient of said first receiving chamber being different from said second physical state of said ingredient of said second receiving chamber.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule further comprises a base and a corresponding cap, wherein said cap is configured to provide a sealing relationship when engaging said base.

10 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said base and said cap are formed having different colors.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said sealing relationship between said base and corresponding cap comprises no dead volume space within said capsule.

15 In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber comprises a pharmaceutical.

In another embodiment of the present invention, there is a multi-compartment capsule  
20 as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber is selected from the group consisting of a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is a multi-compartment capsule  
25 as defined above, wherein said first physical state of said ingredient in said first receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said second physical state of said ingredient in said second

receiving chamber capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule comprises a time-release coating.

5 In another embodiment of the present invention, there is a multi-compartment capsule as defined in above, wherein said capsule is formed of a material selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phtalated gelatin, succinated gelatin, cellulosephtalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, oleoresin,  
10 polymerisates of acrylic or methacrylic esters, polyvinylacetate-phtalate and mixtures thereof.

In another embodiment of the present invention, there is a multi-compartment capsule as defined above, wherein said capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

In another embodiment of the present invention, there is a multi-compartment capsule  
15 as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and mixtures thereof.

In another embodiment of the present invention, there is an encapsulation process for forming a multi-compartment capsule, said process comprising the steps of providing a  
20 primary capsule having a base and a cap; providing a secondary capsule having a base and a cap; introducing at least one ingredient having a first physical state into said secondary capsule; positioning said cap of said secondary capsule in sealing relationship with said base; introducing at least one ingredient having a second physical state into said primary capsule, wherein said first physical state of said ingredient of said secondary capsule is different from  
25 said second physical state of said ingredient of said primary capsule; introducing said secondary capsule into said base of said primary capsule; and positioning said cap of said primary capsule in sealing relationship with said base.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of reducing dead volume space within said  
30 primary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the step of introducing a filling material into said cap of said primary capsule to reduce dead volume space.

5 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said cap of said primary capsule comprises a configuration sufficient for reducing dead volume space within the primary capsule.

10 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient introduced into said primary capsule is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said primary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

15 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said secondary capsule is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

20 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said secondary capsule is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said primary capsule comprises a pharmaceutical and said ingredient in said secondary capsule is selected from the group consisting of a pharmaceutical.

25 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said primary capsule comprises a pharmaceutical and said ingredient in said secondary capsule is selected from the group consisting of a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient introduced into said primary capsule is the same as said ingredient introduced into said secondary capsule.

5 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said time-release coating of said secondary capsule is different from said time-release coating of said primary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the steps of providing a tertiary capsule having a base and a cap; introducing at least one ingredient having a third physical state into said tertiary  
10 capsule; positioning said cap of said secondary capsule in sealing relationship with said base; and introducing said tertiary capsule into said base of said secondary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said tertiary capsule is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary  
15 supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said tertiary capsule comprises a physical state selected from the group consisting of a solid, a liquid, a gas and a dispersion.

20 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said tertiary capsule comprises a time-release coating.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and combinations thereof.

25 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said primary and secondary capsules are formed having different colors.

In another embodiment of the present invention, there is an encapsulation process for forming a multi-compartment capsule, said process comprising the steps of providing a

capsule comprising a cap, a base configured having a longitudinally extending body including a length and at least one dividing wall formed along said length of said extending body, said dividing wall adapted to form a first receiving chamber and a second receiving chamber; introducing at least one ingredient having a first physical state into said second  
5 receiving chamber; introducing at least one ingredient having a second physical state into said first receiving chamber, wherein said first physical state of said ingredient of said second receiving chamber being different from said second physical state of said ingredient of said first receiving chamber; and positioning said cap in sealing relationship with said base.

In another embodiment of the present invention, there is an encapsulation process as  
10 defined above, further comprising the step of reducing dead volume space within said primary capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said filling material is selected from the group consisting of gelatin, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum,  
15 carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, polyvinylacetate-phthalate, polymerisates of acrylic or methacrylic esters and combinations thereof.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said cap comprises a configuration sufficient for reducing dead  
20 volume space within said capsule.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said first receiving chamber is selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as  
25 defined above, wherein said physical state of said ingredient in said receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said second receiving chamber is selected from the

group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient in said second receiving  
5 chamber is selected from the group consisting of a solid, a liquid, a gas and a dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient in said first receiving chamber comprises a pharmaceutical and said ingredient in said second receiving chamber is selected from the group consisting of a pharmaceutical.

10 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said time-release coating of said second receiving chamber is different from said time-release coating of said first receiving chamber.

In another embodiment of the present invention, there is an encapsulation process as defined above, further comprising the steps of positioning a second dividing wall along said  
15 length of said extending body of said base, said second dividing wall adapted to form a third receiving chamber; and introducing at least one ingredient having a physical state into said third receiving chamber.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said ingredient introduced into said third receiving chamber is  
20 selected from the group consisting of a pharmaceutical, a biotechnical, a nutraceutical, a vitamin, a dietary supplement and a mineral.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said physical state of said ingredient introduced into said third receiving chamber is selected from the group consisting of a solid, a liquid, a gas and a  
25 dispersion.

In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said second dividing wall comprises a time-release coating.



In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said capsule further comprises a soft elastic capsule formed of a material selected from the group consisting of glycerin and sorbitol.

5 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said lubricant is selected from the group consisting of aluminiumstearate, calciumstearate, magnesiumstearate, tinstearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones and combinations thereof.

10 In another embodiment of the present invention, there is an encapsulation process as defined above, wherein said base and said cap of said capsule are formed having different colors.

#### BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects and features of the present invention will become more fully apparent from the following description and appended claims, taken in  
15 conjunction with the accompanying drawings. Understanding that these drawings depict only typical embodiments of the invention and are, therefore, not to be considered limiting of its scope, the invention will be described with additional specificity and detail through use of the accompanying drawings in which:

Figure 1 is a flow diagram illustrating one presently preferred embodiment of a  
20 process of the present invention comprising the steps of introducing at least one active ingredient or medicament having a solid physical state into a secondary capsule and introducing the secondary capsule into a primary capsule further including at least one active ingredient or medicament having a liquid physical state;

Figure 2 is a cross-sectional view illustrating another presently preferred embodiment  
25 of a multi-compartment capsule of the present invention wherein a primary capsule houses a secondary capsule and a secondary capsule houses a tertiary capsule, wherein each of the capsules include one or more active ingredients or medicaments and the active ingredient(s) introduced into at least two of the capsules comprise different physical states;

Figure 3 is a perspective view illustrating yet another presently preferred embodiment of a multi-compartment capsule comprising a base, a cap and a dividing wall positioned between the base and the cap, wherein the dividing wall facilitates the formation of at least two, independent receiving chambers for receiving one or more active ingredients or medicaments having different physical states;

Figure 4 is a cross-sectional view of the multi-compartment capsule shown in Figure 3 wherein the base, the dividing wall defining the two receiving chambers and the cap are assembled to form a capsule of the present invention and wherein one or more active ingredients or medicaments having different physical states are introduced into the receiving chambers;

Figure 5 is a perspective view illustrating an alternate presently preferred embodiment of a multi-compartment capsule of the present invention having a primary capsule comprising a capsular base configured with a longitudinally extending body, a corresponding cap and a series of dividing walls disposed in spaced apart relationship along the length of the longitudinally extending body of the base, wherein the dividing walls define a plurality of independent receiving chambers having an internal periphery sufficient for introducing one or more active ingredients or medicaments having different physical states therein and for introducing a secondary capsule, having one or more active ingredients contained therein, within at least one of said receiving chambers;

Figure 6 is a cross-sectional view of the multi-compartment capsule shown in Figure 5 wherein the base and the cap are assembled to form a single dosage capsule having a series of dividing walls that define a plurality of chambers for receiving one or more active ingredients or medicaments, wherein the active ingredient(s) in at least two of the receiving chambers comprise different physical states;

Figure 7 is a perspective view illustrating yet another presently preferred embodiment of a multi-compartment capsule of the present invention having a primary capsule comprising a capsular base configured with a longitudinally extending body, a corresponding cap and a series of dividing walls disposed in spaced apart relationship, both vertically and horizontally,

along the length of the longitudinally extending body of the base, wherein the dividing walls define a plurality of independent receiving chambers having an internal periphery sufficient for introducing one or more active ingredients or medicaments having different physical states therein;

5           Figure 8 is a perspective view illustrating an alternate preferred embodiment of the multi-compartment capsule shown in Figure 7, wherein the multi-compartment capsule includes a primary capsule comprising a capsular base configured with a longitudinally extending body, a corresponding cap and a series of dividing walls disposed in spaced apart relationship, both vertically and horizontally, along the length of the longitudinally extending  
10   body of the base, wherein the dividing walls define a plurality of independent receiving chambers having an internal periphery sufficient for introducing one or more active ingredients or medicaments having different physical states therein and for introducing a secondary capsule, having one or more active ingredients contained therein, within at least one of said receiving chambers;

15           Figure 9 is a perspective view illustrating yet another presently preferred embodiment of a multi-compartment capsule of the present invention wherein the multi-compartment capsule shown in Figure 7 is introduced within the internal periphery of a receiving chamber of a primary capsule having one or more active ingredients also contained therein;

          Figure 10 is a cross-sectional view illustrating a presently preferred embodiment of a  
20   multi-compartment capsule of the present invention including a secondary capsule having one or more active ingredients or medicaments selectively introduced into the internal periphery of a primary capsule having one or more active ingredients or medicaments, wherein the active ingredient(s) introduced into the primary capsule comprises a physical state ( *e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active  
25   ingredient(s) introduced into the internal periphery of the secondary capsule, the primary capsule further comprising a cap having a generally U-shaped configuration adapted to provide a sealing relationship when engaging the corresponding base, thereby reducing dead space volume in the internal periphery of the receiving chamber of the base;

Figure 11 is a perspective view illustrating yet another presently preferred embodiment of a multi-compartment capsule of the present invention including a secondary capsule having one or more active ingredients or medicaments and having a size and shape sufficient for being selectively introduced into the internal periphery of a primary capsule having one or more active ingredients or medicaments, wherein the active ingredient(s) introduced into the primary capsule comprises a physical state ( *e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active ingredient(s) introduced into the internal periphery of the secondary capsule, the primary capsule further comprising a filling material introduced into the internal periphery of the cap having a general conical configuration and adapted to provide a sealing relationship when engaging the corresponding base, thereby reducing dead space volume in the internal periphery of the receiving chamber of the base;

Figure 12 is a cross-sectional view of the multi-compartment capsule shown in Figure 11 wherein a sufficient amount of filling material is introduced into the internal periphery of the cap, thereby functioning to eliminate or significantly reduce the dead space volume in the receiving chamber of the primary capsule; and

Figure 13 is a cross-sectional view illustrating an alternate presently preferred embodiment of a multi-compartment capsule of the present invention comprising a tertiary capsule having one or more active ingredients or medicaments and having a size a shape sufficient for being introduced into at least a portion of the internal periphery of the receiving chamber of a secondary capsule having one or more active ingredients or medicaments also introduced therein, the size and shape of the secondary capsule sufficient for being selectively introduced into the internal periphery of a primary capsule having one or more active ingredients or medicaments, wherein the active ingredient(s) introduced into the primary capsule comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active ingredient(s) introduced into the receiving chambers of the secondary and tertiary capsules, the primary capsule further comprising a filling material introduced into the internal periphery of the cap having a general conical configuration and

adapted to provide a sealing relationship when engaging the corresponding base, thereby reducing dead space volume in the internal periphery of the receiving chamber of the base of the primary capsule.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

It will be readily understood that the components of the present invention, as generally described and illustrated in the Figures herein, could be arranged and designed in a wide variety of different configurations and process steps. Those of ordinary skill in the art will, of course, appreciate that various modifications to the details herein may easily be made without departing from the essential characteristics of the invention, as described. Thus, the following more detailed description of the embodiments of apparatus and methods of the present invention, as represented in Figures 1 through 13, is not intended to limit the scope of the invention, as claimed, but it is merely representative of the presently preferred embodiments of the invention.

The presently preferred embodiments of the invention will be best understood by reference to the drawings, wherein like parts are designated by like numerals throughout.

One presently preferred embodiment of the present invention, designated generally at 10, is best illustrated in Figure 1. As shown, a multi-compartment capsule 10 is illustrated comprising a primary capsule 11 and a secondary capsule 20 selectively introduced within at least a portion of an internal periphery of the primary capsule. The secondary capsule 20 includes a base 24, a corresponding cap 22 and a receiving chamber 28 formed between the base and cap. The receiving chamber 28 is configured having an internal periphery sufficient for receiving at least one active ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. In similar structural design, the primary capsule 11 may be formed having a base 14, a corresponding cap 12 and a receiving chamber 18 formed between the base and cap. The receiving chamber 18 of the primary capsule 11 is preferably formed having an internal periphery sufficient for receiving the secondary capsule 20, together with at least one active

ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein.

Still referring to Figure 1, one presently preferred embodiment of an encapsulation process for forming a multi-compartment capsule 10 is comprising the steps of: (1) providing  
5 a primary capsule 11 having a base 14, a corresponding cap 12 and a receiving chamber 18;  
(2) providing a secondary capsule 20 having a base 24, a corresponding cap 22 and a receiving chamber 28; (3) introducing at least one ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a first physical state (*e.g.*, solid, liquid, gas or dispersion) into at  
10 least a portion of the receiving chamber 28 of the secondary capsule 20 and selectively positioning the cap 22 in sealing relationship with the base 24; (4) introducing at least one ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a second physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber 18 of the primary  
15 capsule 11, wherein the first physical state of the ingredient(s) in the secondary capsule is different from the second physical state of the ingredient(s) in the primary capsule; and (5) introducing the secondary capsule 20 into at least a portion of the receiving chamber 18 of the primary capsule 11 and selectively positioning the cap 12 in sealing relationship with the base 14 to form a single dosage multi-compartment capsule.

20 As shown, a solid is selectively introduced within at least a portion of the internal periphery of the receiving chamber 28 of the secondary capsule 20 and a liquid is selectively introduced within at least a portion of the internal periphery of the receiving chamber 18 of the primary capsule 11. Although the ingredient(s) introduced into the receiving chamber 18 of the primary capsule 11 may be the same or different from the ingredient(s) introduced into  
25 the receiving chamber 28 of the secondary capsule, the active ingredient(s) in the primary capsule 11 have a physical state (*i.e.*, solid, liquid, gas or dispersion) that varies from the physical state of the active ingredient(s) in the secondary capsule 20. Accordingly, those skilled in the art will readily recognize other possible modifications and adaptations relative

to the contemplated variations in physical states of the active ingredient(s) selectively positionable within the receiving chambers 18, 28 of the primary and secondary capsules, respectively, which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figure 2, an alternate presently preferred embodiment of a multi-compartment capsule 110 is shown comprising a primary capsule 111, a secondary capsule 120 and a tertiary capsule 130. The tertiary capsule 130 includes a base 134, a corresponding cap 132 and a receiving chamber 138 formed between the base and cap. The receiving chamber 138 is preferably formed having an internal periphery sufficient for receiving at least one active ingredient or medicament (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof). Structurally, the tertiary capsule 130 is configured having a size sufficient for being selectively introduced within at least a portion of an internal periphery of a receiving chamber 128 defined between a base 124 and a corresponding cap 122 of the secondary capsule 120. One or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) may be introduced into at least a portion of the receiving chamber 128 of the secondary capsule 120, together with the introduction of the tertiary capsule 130 comprising its active ingredient(s). The secondary capsule 120 having its active ingredient(s) and housing the tertiary capsule 130 with its active ingredient(s) may then be selectively introduced within at least a portion of an internal periphery of a receiving chamber 118 of the primary capsule 111 defined between a base 124 and a corresponding cap 122. Preferably, the receiving chamber 118 of the primary capsule 111 may also include one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) introduced therein.

Still referring to Figure 2, another presently preferred embodiment of an encapsulation process for forming a multi-compartment capsule 110 may comprise the steps of: (1) providing a primary capsule 111 having a base 114, a corresponding cap 112 and a receiving chamber 118 defined between the base and cap; (2) providing a secondary capsule 120 having a base 124, a corresponding cap 122 and a receiving chamber 128 defined between the base and cap; (3) providing a tertiary capsule 130 having a base 134, a corresponding cap 132 and a receiving chamber 138 defined between the base and cap; (4) introducing at least one ingredient (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a first physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber 138 of the tertiary capsule 130 and selectively positioning the cap 132 in sealing relationship with the base 134; (5) introducing at least one ingredient (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a second physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber 128 of the secondary capsule 120, wherein the first physical state of the ingredient(s) in the tertiary capsule 130 are the same as the second physical state of the ingredient(s) in the secondary capsule 120; (6) introducing the tertiary capsule 130 into at least a portion of the receiving chamber 218 of the secondary capsule 120 and selectively positioning the cap 122 in sealing relationship with the base 124; (7) introducing at least one ingredient (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a third physical state (*e.g.*, solid, liquid, gas or dispersion) into at least a portion of the receiving chamber 118 of the primary capsule 111, wherein the third physical state of the ingredient(s) in the primary capsule are different from the first and second physical states of the ingredient(s) in the tertiary capsule 130 and the secondary capsule 120, respectively; and (8) introducing the secondary capsule 120 into at least a portion of the receiving chamber 118 of the primary capsule 111 and selectively positioning the cap 112 in sealing relationship with the base 114 to form a single dosage multi-compartment capsule.



In the presently preferred embodiment illustrated in Figure 2, a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 118 of the primary capsule 111, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 128 of the secondary capsule 120 and a solid may be selectively introduced into at least a portion of the receiving chamber 138 of the tertiary capsule 130. Although the ingredient(s) selectively introduced into the receiving chambers 118, 128, 138 of the primary, secondary and tertiary capsules 111, 120, 130, respectively, may be the same or different, the active ingredient(s) in at least two of the receiving chambers comprise at least two different physical states (*e.g.*, solid, liquid, gas or dispersion). It is further contemplated herein as an alternate embodiment that the active ingredient(s) introduced in the receiving chamber 118 of the primary capsule 111 comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of the active ingredient(s) contained within the receiving chamber 128 of the secondary capsule 120 which is different from the physical state of the active ingredient(s) contained within the receiving chamber 138 of the tertiary capsule 130. Those skilled in the art will readily recognize other possible modifications and adaptations relative to contemplated variations in physical states of the active ingredient(s) selectively introduced within the receiving chambers 118, 128, 138 of the primary, secondary and tertiary capsules, respectively, which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figures 3 and 4, another presently preferred embodiment of a multi-compartment capsule 210 is shown comprising a base 214, a corresponding cap 212 and a dividing wall 216 positionable between the base and the cap. Structurally, the size, shape and positioning of the dividing wall 216 relative to the base 214 and corresponding cap 212 facilitates the formation of at least two, independent and separate receiving chambers 218a, 218b, each having an internal periphery sufficient for receiving one or more active

ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. As best shown in Figure 4, the dividing wall 216 seats within the internal periphery of both the base 214 and the corresponding cap 212. After introducing one or more active ingredients or medicaments  
5 into receiving chamber 218b and disposing the dividing wall 216 relative thereto, one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) may be introduced into receiving chamber 218a and the cap may be selectively positioned in sealing relationship with the base 214 to form one presently preferred embodiment of the single, dosage  
10 multi-compartment capsule 210. Moreover, the dividing wall 216 may functionally assist in forming a sealing relationship between the base 214 and corresponding cap 212 of the multi-compartment capsule 210, if desired.

In one presently preferred embodiment of the multi-compartment capsule 211 of the present invention, a solid may be selectively introduced into at least a portion of the internal  
15 periphery of the receiving chamber 218a and a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 218b. Although the ingredient(s) introduced into the receiving chamber 218a may be the same or different from the ingredient(s) introduced into the receiving chamber 218, the active ingredient(s) in the first receiving chamber 218a preferably comprise a physical state (*e.g.*, solid, liquid, gas or  
20 dispersion) that is different from the physical state of the active ingredient(s) in the second receiving chamber 218b. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively positionable within the receiving chambers 218a, 218b which are consistent with the spirit and scope of the present  
25 invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figures 5 and 6, another presently preferred embodiment of a multi-compartment capsule, designated as 310, is shown including a primary capsule 311 comprising a capsular base 314 configured having an elongated or longitudinally extending body, a corresponding cap 312 and a plurality of dividing walls 316 selectively disposed  
5 along the length of the longitudinally extending body of the base. Preferably, the structural size, shape and positioning of the dividing walls 316a, 316b, 316c along the length of the elongated body of the base 314 facilitate the formation of a plurality of independent receiving chambers 318a, 318b, 318c, 318d. Each receiving chamber 318a, 318b, 318c, 318d of the primary capsule 311 having an internal periphery sufficient for receiving one or more active  
10 ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein.

As best shown in Figure 6, the dividing walls 316a, 316b, 316c are preferably seated within the internal periphery of the base 314 of the primary capsule 311 and in a spaced apart relationship along the length of the longitudinally extending body and form four independent  
15 receiving chambers 318a, 318b, 318c, 318d. In one presently preferred embodiment of the multi-compartment capsule 310 of the present invention, each of the receiving chambers 318a, 318b, 318c comprises at least one active ingredient or medicament having a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of the ingredient(s) in the other receiving chambers.

As illustrated by way of example, and not by way of restriction, a solid may be  
20 selectively introduced into at least a portion of the internal periphery of the receiving chamber 318a, a dispersion may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 318b, a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 318c and a secondary  
25 capsule 320 may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 318d. As contemplated herein, receiving chamber 318d may be further configured having an internal periphery sufficient for receiving a secondary capsule 320, together with at least one active ingredient or medicament therein.

One presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 310 illustrated in Figures 5 and 6, may comprise the steps of: (1) introducing a secondary capsule 320 (*e.g.*, tablet) and one or more active ingredients or medicaments into receiving chamber 318d; (2) selectively positioning dividing wall 316c along the length of the elongated body of the base 314; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 318c; (4) selectively positioning dividing wall 316b along the length of the elongated body of the base 314 in a spaced apart relationship to dividing wall 316c; (5) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 318b; (6) selectively positioning dividing wall 316a along the length of the elongated body of the base 314 in a spaced apart relationship to dividing wall 316b; and (7) selectively positioning the cap 312 in sealing relationship with the base 314 to form a presently preferred embodiment of a single, dosage multi-compartment capsule 310. The dividing wall 316a may also function in the formation of the sealing relationship between the base 314 and the corresponding cap 312, if desired.

Although the ingredient(s) introduced into one of the receiving chambers 318 may be the same ingredient or may be different from the ingredient(s) introduced into the other receiving chambers, the active ingredient(s) in at least two of the receiving chambers 318 preferably comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in one or more of the remaining receiving chambers. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively introduced within the receiving chambers 318 which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of

the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Another presently preferred embodiment of a multi-compartment capsule of the present invention, generally designated as 410 in Figure 7, is shown comprising a capsular  
5 base 414 preferably configured having an elongated or longitudinally extending body, a corresponding cap 412 and a plurality of dividing walls 416 selectively disposed along the length of the longitudinally extending body of the base, both horizontally and vertically. In structural design, the size, shape and positioning of the dividing walls 416a, 416b, 416c, 416d, 416e along the length of the longitudinally extending body of the base 414 facilitate the  
10 formation of a plurality of independent receiving chambers 418.

In one presently preferred embodiment, the dividing walls 416a, 416b, 416c, 416d, 416e are preferably disposed or seated in a spaced apart relationship within the internal periphery of the base 414 of the primary capsule 411 along the length of the longitudinally extending body, whereby forming five (5) independent receiving chambers 418a, 418b, 418c,  
15 418d, 418e. Each receiving chamber 418a, 418b, 418c, 418d, 418e of the primary capsule 411 are preferably configured having an internal periphery dimensionally sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein.

Still referring to Figure 7, one presently preferred embodiment of an encapsulation  
20 process, as defined by the structural configuration of the multi-compartment capsule 410, may comprise the steps of: (1) introducing one or more active ingredients or medicaments into receiving chamber 418e defined by dividing walls 416d, 416e which are vertically disposed along the length of the elongated body of the base 414; (2) introducing one or more active ingredients or medicaments into receiving chamber 418d defined by dividing walls  
25 416c, 416d which are vertically disposed along the length of the elongated body of the base 414; (3) introducing one or more active ingredients or medicaments into receiving chamber 418c defined by dividing walls 416b, 416c which are vertically disposed along the length of the elongated body of the base 414; (4) introducing one or more active ingredients or

medicaments into receiving chamber 418b defined by dividing walls 416b, 416e which are vertically disposed along the length of the elongated body of the base 414; (5) disposing dividing wall 416a along the length of the elongated body of the base 414 perpendicular to the disposition of dividing walls 416b, 416c, 416d, 416e and introducing one or more active ingredients or medicaments into receiving chamber 418a; and (6) selectively positioning the cap 412 in sealing relationship with the base 414 to form one presently preferred embodiment of a single, dosage multi-compartment capsule 410. As appreciated, the dividing wall 416a may also function in the formation of the sealing relationship between the base 414 and the corresponding cap 412, if structurally desired.

As illustrated by way of example, and not by way of restriction, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 418a, a dispersion may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 418b, a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 418c, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 418d and a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 418e.

Although the ingredient(s) introduced into one of the receiving chambers 418 may be the same ingredient or may be different from the ingredient(s) introduced into the other receiving chambers, the active ingredient(s) in at least two of the receiving chambers 418 preferably comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in one or more of the remaining receiving chambers. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively introduced within the receiving chambers 418 which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of

the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figure 8, an alternate presently preferred embodiment of a multi-compartment capsule 510 includes a capsular base 514 preferably configured having an elongated or longitudinally extending body, a corresponding cap 512 and a plurality of dividing walls 516 selectively disposed along the length of the longitudinally extending body of the base, both horizontally and vertically. In structural design, the size, shape and positioning of the dividing walls 516a, 516b, 516c, 516d along the length of the longitudinally extending body of the base 514 facilitate the formation of a plurality of independent receiving chambers 518.

In one presently preferred embodiment, the dividing walls 516a, 516b, 516c, 516d, 516e are preferably disposed or seated in a spaced apart relationship within the internal periphery of the base 514 of the primary capsule 511 along the length of the longitudinally extending body, whereby forming five (5) independent receiving chambers 518a, 518b, 518c, 518d, 518e. Each of the receiving chamber 518a, 518b, 518c, 518d, 518e of the primary capsule 411 are preferably configured having an internal periphery dimensionally sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. Moreover, receiving chamber 518d is formed having an internal periphery sufficient for receiving a secondary capsule 520. The secondary capsule 520 being configured with a base 524, corresponding cap 522 and a dividing wall 526 defining a first receiving chamber 528a and a second receiving chamber 528b. The first receiving chamber 528a is preferably configured having an internal periphery sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) having a first physical state (*e.g.*, solid, liquid, gas or dispersion) therein. Similarly, the second receiving chamber 528b is configured having an internal periphery sufficient for receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement,

mineral or combination thereof) having a second physical state (*e.g.*, solid, liquid, gas or dispersion), wherein the physical state of the ingredient(s) in the second receiving chamber varies from the physical state of the ingredient(s) in the first receiving chamber. As contemplated and disclosed hereinabove, after the ingredients are introduced into the  
5 respective receiving chambers 528a, 528b, the cap 522 may be positioned in sealing relationship with the base 524 of the secondary capsule 520.

Still referring to Figure 8, as illustrated by way of example, and not by way of restriction, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 528a and a liquid may be selectively introduced into at  
10 least a portion of the internal periphery of the receiving chamber 528b. Although the ingredient(s) introduced into one of the receiving chamber 528a may be the same ingredient or may be different from the ingredient(s) introduced into receiving chamber 528b, the active ingredient(s) in the first receiving chamber 528a comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in  
15 receiving chambers 528b. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively introduced within the receiving chambers 528 which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed  
20 as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles

One presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 510, may comprise the steps of: (1) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical,  
25 biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 518e defined by dividing walls 516d, 516e which are disposed vertically along the length of the elongated body of the base 514; (2) introducing a secondary capsule 520 into receiving chamber 518d defined by dividing walls 516c, 516d which are disposed



vertically along the length of the elongated body of the base 514; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 518c defined by dividing walls 516b, 516c which are disposed vertically along the length of the elongated body of the base 514; (4) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 518b defined by dividing walls 516b, 516e which are disposed vertically along the length of the elongated body of the base 514; (5) disposing dividing wall 516a along the length of the elongated body of the base 514 perpendicular to the disposition of dividing walls 516b, 516c, 516d, 516e and introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 518a; and (6) selectively positioning the cap 512 in sealing relationship with the base 514 to form one presently preferred embodiment of a single, dosage multi-compartment capsule 510. As appreciated, the dividing wall 516a may also function in the formation of the sealing relationship between the base 514 and the corresponding cap 512, if structurally desired.

As illustrated by way of example, and not by way of limitation, a solid may be selectively introduced into at least a portion of the internal periphery of receiving chamber 518a, a dispersion may be selectively introduced into at least a portion of the internal periphery of receiving chamber 518b, a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 518c and a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 518e. Although the ingredient(s) introduced into one of the receiving chambers 518 may be the same ingredient or may be different from the ingredient(s) introduced into the other receiving chambers, the active ingredient(s) in at least two of the receiving chambers 518 preferably comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in one or more of the remaining receiving

chambers. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively introduced within the receiving chambers 518 which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figure 9, yet another presently preferred embodiment of a multi-compartment capsule of the present invention, generally designated as 610, is shown comprising a primary capsule 611 and a secondary capsule 620 selectively positionable within at least a portion of an internal periphery of the primary capsule. The primary capsule 611 having a receiving chamber 618 preferably formed having an internal periphery sufficient for receiving the secondary capsule 620, together with one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein. The secondary capsule 620 comprising a capsular base 624 preferably configured having an elongated or longitudinally extending body, a corresponding cap 622 and a plurality of dividing walls 626 selectively disposed along the length of the longitudinally extending body of the base, both horizontally and vertically. In structural design, the size, shape and positioning of the dividing walls 626a, 626b, 626c, 626d along the length of the longitudinally extending body of the base 624 facilitate the formation of a plurality of independent receiving chambers 628.

In one presently preferred embodiment, the dividing walls 626a, 626b, 626c, 626d, 626e are preferably disposed or seated in a spaced apart relationship within the internal periphery of the base 624 of the secondary capsule 620 along the length of the longitudinally extending body, whereby forming five (5) independent receiving chambers 628a, 628b, 628c, 628d, 628e. Each receiving chamber 628a, 628b, 628c, 628d, 628e of the secondary capsule 620 are preferably configured having an internal periphery dimensionally sufficient for

receiving one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) therein.

One presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 610, may comprise the steps of: (1) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 628e defined by dividing walls 626d, 626e which are vertically disposed along the length of the elongated body of the base 624; (2) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 628d defined by dividing walls 626c, 626d which are vertically disposed along the length of the elongated body of the base 624; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 628c defined by dividing walls 626b, 626c which are vertically disposed along the length of the elongated body of the base 624; (4) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 628b defined by dividing walls 626b, 626e which are vertically disposed along the length of the elongated body of the base 624; (5) disposing dividing wall 626a along the length of the elongated body of the base 624 perpendicular to the disposition of dividing walls 626b, 626c, 626d, 626e and introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 628a; (6) selectively positioning the cap 622 in sealing relationship with the base 624 of the secondary capsule 620; (7) introducing the secondary capsule 620 and one or more ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into the receiving chamber 618 of the primary capsule 611; and (8) selectively positioning the cap 612 in sealing relationship with the base 614 of the primary

capsule 611 to form one presently preferred embodiment of a single, dosage multi-compartment capsule 610.

As illustrated by way of example, and not by way of restriction, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 628a, a dispersion may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 628b, a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 628c, a solid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 628d and a liquid may be selectively introduced into at least a portion of the internal periphery of the receiving chamber 628e of the secondary capsule 620. In addition, a gas may be introduced into at least a portion of the internal periphery of the receiving chamber 618 of the primary capsule 611.

Although the ingredient(s) introduced into one of the receiving chambers 618, 628 of the primary and secondary capsules, respectively, may be the same ingredient or may be different from the ingredient(s) introduced into the other receiving chambers, the active ingredient(s) in at least two of the receiving chambers 618, 628 preferably comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) that is different from the physical state of the active ingredient(s) in one or more of the remaining receiving chambers. Those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states (*e.g.*, solid, liquid, gas and dispersion) of the active ingredient(s) selectively introduced within the receiving chambers 618, 628 which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Another presently preferred embodiment of a multi-compartment capsule of the present invention, generally designated as 710 in Figure 10, is shown comprising a secondary capsule 720 including one or more active ingredients or medicaments (*e.g.*, pharmaceutical,

biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) within at least a portion of the internal periphery of a receiving chamber 728 and having a size and shape sufficient for being selectively introduced within at least a portion of the internal periphery of a receiving chamber 718 of a primary capsule 711. The primary capsule 5 711 also includes one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) introduced within the internal periphery of the receiving chamber 718, wherein the active ingredient(s) introduced into the primary capsule comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active ingredient(s) 10 introduced into the internal periphery of the secondary capsule. In structural design, the primary capsule 711 further comprises a cap 712 having a generally U-shaped configuration adapted to provide a sealing relationship when engaging the corresponding base 714, thereby reducing dead space volume in the internal periphery of the receiving chamber 718 of the base. In this regard, the configuration of the cap 712 generally eliminates or substantially 15 reduces the potential dead space volume within the internal periphery of the receiving chamber 718, thus functionally negating the opportunity for reaction between an air bubble and the active ingredient(s) introduced into the base 714 of the primary capsule 711.

One presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 710, may include the steps of: (1) 20 introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into receiving chamber 728; (2) selectively positioning the cap 722 in sealing relationship with the base 724 of the secondary capsule 720; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, 25 mineral or combination thereof), together with the secondary capsule 720, into the receiving chamber 718 of the primary capsule 711; and (4) selectively positioning the cap 712 having a general U-shaped configuration in sealing relationship with the base 714 of the primary capsule 711 to form a presently preferred embodiment of a single, dosage multi-compartment

capsule 710, wherein eliminating or substantially reducing dead space volume within the internal periphery of the receiving chamber 718.

A solid is selectively introduced within at least a portion of the internal periphery of the receiving chamber 728 of the secondary capsule 720 and a liquid is selectively introduced within at least a portion of the internal periphery of the receiving chamber 718 of the primary capsule 711. Although the ingredient(s) introduced into the receiving chamber 718 of the primary capsule 711 may be the same or different from the ingredient(s) introduced into the receiving chamber 728 of the secondary capsule 720, the active ingredient(s) in the primary capsule have a physical state (*i.e.*, solid, liquid, gas or dispersion) that various from the physical state of the active ingredient(s) in the secondary capsule. Accordingly, those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states of the active ingredient(s) selectively introduced within the receiving chambers 718, 728 of the primary and secondary capsules 711, 720, respectively, which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figures 11 and 12, yet another presently preferred embodiment of a multi-compartment capsule 810 of the present invention is shown comprising a secondary capsule 820 including one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) within at least a portion of the internal periphery of a receiving chamber 828. The secondary capsule 820 being preferably formed having a size and shape sufficient for being selectively introduced within at least a portion of the internal periphery of a receiving chamber 818 of a primary capsule 811. Similarly, the primary capsule 811 includes one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) introduced within the internal periphery of the receiving chamber 818, together with the secondary capsule 820, wherein the active

ingredient(s) introduced into the primary capsule comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active ingredient(s) introduced into the internal periphery of the secondary capsule 820.

In preferred structural design, the primary capsule 811 comprises a cap 812 having a general cylindrical configuration adapted to provide a sealing relationship when engaging the corresponding base 814 to form a single dosage, multi-compartment capsule 810. An amount of filling material 840 may be introduced into the internal periphery of the cap 812 to fill, either partially or completely, the inner volume of the cap, thereby reducing the dead space volume in the internal periphery of the receiving chamber 818 of the base. In this regard, the configuration of the addition of the filler material 840 relative to the internal periphery of the cap 812 generally eliminates or substantially reduces the potential dead space volume within the internal periphery of the receiving chamber 818, thus functionally negating the potential for a reaction between an air bubble and the active ingredient(s) introduced into the base 814 of the primary capsule 811.

Preferably, the filling material 840 introduced into at least a portion of the internal periphery of the cap 812 may include a hydrophilic polymer, such as gelatin. It will be readily appreciated by those skilled in the art that other filling materials may be used, such as, for example, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose, (HPMC), oleoresin, polyvinylacetate-phthalate, polymerisates of acrylic or methacrylic esters, and mixtures thereof, or the like, and/or combinations thereof. In other presently preferred embodiments of the present invention, the filling material 840 may include the introduction of an inert compound, for example, nitrogen gas into at least a portion of the internal periphery of the cap 811. Based on the principals of eliminating or reducing the volume dead space in multi-compartment capsules disclosed herein, those skilled in the art will readily recognize other possible modifications and combinations which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the examples provided

herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or process for implementing those principles.

The filling material 840 introduced within at least a portion of the internal periphery of the cap 812 of the primary capsule 811 is generally intended to promote a binding contact with at least a portion of the cap 822 of the secondary capsule 820, thereby seating at least a portion of the secondary capsule within the cap of the primary capsule and forming a molded appearance. As appreciated, the introduction of the filling material 840 into the cap 812 of the primary capsule 811 prior to the joining and sealing process may prevent the opportunity for a reaction between an air bubble and the active medicament(s) within the receiving chamber 818 of the primary capsule, while preserving the overall rounded shape of the multi-compartment capsule 910 for ease of swallowing by a consumer.

As best illustrated in Figure 12, one presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 810, may include the steps of: (1) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin; dietary supplement, mineral or combination thereof) into at least a portion of the receiving chamber 828; (2) selectively positioning the cap 822 in sealing relationship with the base 824 of the secondary capsule 820; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof), together with the secondary capsule 820, into at least a portion of the receiving chamber 818 of the primary capsule 811; (4) introducing a filling material 840 into at least a portion of the internal periphery of the cap 812 (*i.e.*, filling the cap); and (5) selectively positioning the cap 812 having a general conical configuration in sealing relationship with the base 814 of the primary capsule 811 to form one presently preferred embodiment of a single, dosage multi-compartment capsule 810, wherein eliminating or substantially reducing dead space volume within the internal periphery of the cap 812 and the receiving chamber 818, respectively.



A solid may be selectively introduced within at least a portion of the internal periphery of the receiving chamber 828 of the secondary capsule 820 and a liquid may be selectively introduced within at least a portion of the internal periphery of the receiving chamber 818 of the primary capsule 811. Although the ingredient(s) introduced into the receiving chamber 818 of the primary capsule 811 may be the same or different from the ingredient(s) introduced into the receiving chamber 828 of the secondary capsule 820, the active ingredient(s) in the primary capsule have a physical state (*i.e.*, solid, liquid, gas or dispersion) that various from the physical state of the active ingredient(s) in the secondary capsule. Accordingly, those skilled in the art will readily recognize other possible modifications and adaptations relative to the contemplated variations in physical states of the active ingredient(s) selectively introduced within the receiving chambers 818, 828 of the primary and secondary capsules 811, 820, respectively, which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

Referring now to Figure 13, another presently preferred embodiment of a multi-compartment capsule, generally designated at 910, is shown comprising a tertiary capsule 930 including one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) within at least a portion of the internal periphery of a receiving chamber 938 and having a size and shape sufficient for being introduced into the internal periphery of a receiving chamber 928 of a secondary capsule 920. The secondary capsule 920 having one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) introduced within at least a portion of the internal periphery of a receiving chamber 928, together with the tertiary capsule 930. The secondary capsule 920 preferably formed having a size and shape sufficient for being selectively introduced within at least a portion of the internal periphery of a receiving chamber 918 of a primary capsule 911. Similarly, the primary capsule 911 may

include one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) introduced within the internal periphery of the receiving chamber 818, together with the secondary capsule 920 which houses the tertiary capsule 930. In one presently preferred embodiment, the active ingredient(s) introduced into the secondary capsule 920 comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) which differs from the physical state of the active ingredient(s) introduced into the internal periphery of the primary capsule 911 and the internal periphery of the tertiary capsule 930.

In preferred structural design, the primary capsule 911 comprises a cap 912 having a general cylindrical configuration adapted to provide a sealing relationship when engaging the corresponding base 914 to form a single dosage, multi-compartment capsule 910. An amount of filling material 940 may be introduced into at least a portion of the internal periphery of the cap 912 to fill, either partially or completely, the inner volume of the cap, thereby reducing the dead space volume in the cap and the internal periphery of the receiving chamber 918 of the base. In this regard, the configuration of the addition of the filler material 940 relative to the internal periphery of the cap 912 may generally eliminate or substantially reduce the potential dead space volume within the internal periphery of the receiving chamber 918, thus functionally negating the potential for a reaction between an air bubble and the active ingredient(s) introduced into the base 914 of the primary capsule 911.

Preferably, the filling material 940 introduced into at least a portion of the internal periphery of the cap 912 may include a hydrophilic polymer, such as gelatin. It will be readily appreciated by those skilled in the art that other filling materials may be used, such as, for example, starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methylcellulose, oleoresin, polyvinylacetate-phthalate, polymerisates of acrylic or methacrylic esters, and mixtures thereof, or the like, and/or combinations thereof. In other presently preferred embodiments of the present invention, the filling material 840 may include the introduction of an inert

compound, for example, nitrogen gas into at least a portion of the internal periphery of the cap 912. Based on the principals of eliminating or reducing the volume dead space in multi-compartment capsules disclosed herein, those skilled in the art will readily recognize other possible modifications and combinations which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or process for implementing those principles.

The filling material 940 introduced within at least a portion of the internal periphery of the cap 912 of the primary capsule 911 is generally intended to promote a binding contact with at least a portion of the cap 922 of the secondary capsule 920, thereby seating at least a portion of the secondary capsule within the cap of the primary capsule and forming a molded appearance. As appreciated, the introduction of the filling material 940 into the cap 912 of the primary capsule 911 prior to the joining and sealing process tends to prevent the opportunity for a reaction between an air bubble and the active medicament(s) within the receiving chamber 918 of the primary capsule, while preserving the overall rounded shape of the multi-compartment capsule 910 for ease of swallowing by a consumer.

As best illustrated in Figure 13, one presently preferred embodiment of an encapsulation process, as defined by the structural configuration of the multi-compartment capsule 910, may include the steps of: (1) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof) into at least a portion of the receiving chamber 938 of a tertiary capsule 930; (2) selectively positioning the cap 932 in sealing relationship with the base 934 of the tertiary capsule 930; (3) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof), together with the tertiary capsule 930, into at least a portion of the receiving chamber 928 of the secondary capsule 920; (4) selectively positioning the cap 922 in sealing relationship with the base 924 of the secondary capsule 920; (5) introducing one or more active ingredients or medicaments (*e.g.*, pharmaceutical,

biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof), together with the secondary capsule 920, into at least a portion of the receiving chamber 918 of the primary capsule 911; (6) introducing a filling material 940 into at least a portion of the internal periphery of the cap 912 (*i.e.*, preferably filling the cap); and (7) selectively  
5 positioning the cap 912 having a general conical configuration in seating relationship with at least a portion of the secondary capsule 920 and sealing the base 914 of the primary capsule 911 to form one presently preferred embodiment of a single, dosage multi-compartment capsule 910, wherein eliminating or substantially reducing dead space volume within the internal periphery of the cap 912 and the receiving chamber 918, respectively.

10 A solid may be introduced within at least a portion of the internal periphery of the receiving chamber 938 of the tertiary capsule 930, a liquid may be introduced into at least a portion of the internal periphery of the secondary capsule 920 and a solid may be selectively introduced within at least a portion of the internal periphery of the receiving chamber 918 of the primary capsule 911. Although the ingredient(s) introduced into the receiving chambers  
15 918, 928, 938 of the primary, secondary and tertiary capsules 911, 920, 930, respectively, may be the same or different from the ingredient(s) introduced into the other receiving chambers, the active ingredient(s) in at least two of the receiving chambers 918, 928, 938 have different physical states (*i.e.*, solid, liquid, gas or dispersion). Those skilled in the art will readily recognize other possible modifications and adaptations relative to the  
20 contemplated variations in physical states of the active ingredient(s) selectively introduced within the receiving chambers 918, 928, 938 of the primary, secondary and tertiary capsules 911, 920, 930, respectively, which are consistent with the spirit and scope of the present invention. It is intended, therefore, that the figures and examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular  
25 structure or method for implementing those principles.

Generally referring to Figures 1-13, the component parts of the presently preferred embodiments of the multi-compartment capsules (*i.e.*, capsular base, corresponding cap and dividing walls) of the present invention may comprise a hydrophilic polymer, such as gelatin

(marine or animal based product). Other suitable materials forming the capsules may include starch, casein, chitosan, soya bean protein, safflower protein, alginates, gellan gum, carrageenan, xanthan gum, phthalated gelatin, succinated gelatin, cellulosephthalate-acetate, polyvinylacetate, hydroxypropyl methyl cellulose (HPMC), oleoresins, polyvinylacetate-phthalate, polymerisates of acrylic or methacrylic esters, and mixtures thereof, or the like, and/or combinations thereof. The material comprising the capsular components may further include between about 0% to 40% of pharmaceutically acceptable plasticizers based upon the weight of the hydrophilic polymer. Plasticizers that may be employed include, for example and not by way of limitation, polyethylene glycol, glycerol, sorbitol, dioctyl-sodium sulfosuccinate, triethyl citrate, tributyl citrate, 1,2-propyleneglycol, mono-acetates, di-acetates, or tri-acetates of glycerol, mixtures thereof, or the like, and/or combinations thereof. As appreciated, plasticizers may also be used in the development of a soft elastic shell, often referred to as a soft gelatin capsule or "soft gel"capsule, for a primary capsule, a secondary capsule and/or a tertiary capsule.

The capsular shell material may contain pharmaceutically acceptable lubricants in the range of about 0% to 10%, based upon the weight of the hydrophilic polymer. Lubricants that may be used include, for example and not by way of limitation, aluminum stearate, calcium stearate, magnesium stearate, tin stearate, talc, sodium lauryl sulfate, lecithins, mineral oils, stearic acid, silicones, mixtures thereof, or the like, and/or combinations thereof.

One presently preferred embodiment of the multi-compartmental capsules of the present invention (*e.g.*, primary capsule, secondary capsule, tertiary capsule, etc.) may include, for example, LICAPS® capsules (for poorly soluble compounds), VCAPS™ capsules (made from cellulosic raw materials), CONI-SNAP® capsules and PRESS-FIT® capsules which are all presently manufactured by Capsugel, a subsidiary of Pfizer, Inc.

In one presently preferred embodiment of an encapsulation process, the primary capsule may be kept under conditions of low humidity within a filling machine during the contemplated steps of rectifying and assembling. In certain embodiments, the primary capsule may contain moisture content in the range of approximately 0% to 6% of the total

weight. Similarly, a secondary capsule, a tertiary capsule, etc. may be processed in the same manner as the primary capsule relative to conditions of low humidity during the steps of rectifying and assembling. As contemplated herein, a moisture content of approximately 0% to 3% by weight is preferable. However, capsules having a higher moisture content than those stated herein are certainly not outside the spirit and scope of the present invention.

As illustrated in Figures 1-9 and 11-13, the shape of the base and corresponding cap of the capsules (*e.g.*, primary, secondary, tertiary, etc.) of the presently preferred embodiments of the multi-compartment capsules are configured having a general cylindrical shape which defines a diameter and length sufficient for the introduction of an internal smaller capsule or one or more dividing walls along the length of the capsular base. It is apparent that other geometrical configurations of the cap are likewise suitable and contemplated herein, such as the general U-shaped configuration of the cap shown in Figure 10. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to any particular structure or configuration for implementing those principles.

In one presently preferred embodiment, the clearance between the primary capsule and the secondary capsule introduced within the internal periphery of the primary capsule is preferably greater than +0.2 mm. The clearance between the outer capsular walls of the secondary capsule and the inner capsular walls of the primary capsule (or the tertiary capsule and the secondary capsule) may be in the range of about 0 mm to 0.5 mm, whereas the outer capsular walls of the secondary capsule or tertiary capsule may be in actual contact with the inner capsular walls of the primary capsule or secondary capsule, respectively. As appreciated, in an effort to structural facilitate independent receiving chambers on opposing sides of a dividing wall introduced within the internal periphery of a base of a capsule, the perimeter of the dividing wall preferably engages the inner capsular walls of the capsule to provide a sealing relationship therebetween.

As further contemplated herein, the inner capsular walls of a primary capsule may be treated with an adhesive sufficient to improve engagement between the primary capsule and

the outer capsular walls of a secondary capsule. A suitable technique to apply an adhesive may be by way of spraying the same on the shells and capsules immediately before assembling the same. Suitable adhesives that may be used may include, for example, tackidex, an aqueous gelatin solution, or the like.

5           The primary, secondary or tertiary capsules, in accordance with the present invention, may be formed having the same or different colors. Moreover, the base and corresponding cap of a single capsule may be formed having different colors in an effort to enhance the aesthetics of the capsule to the consumer. In one presently preferred embodiment of a multi-compartment capsule of the present invention, the dosage may be banded, sealed or  
10 easily dividable in a contact area of the primary and secondary capsules or the sealing band may be color-coded to assist in branding, if desired.

          It is further contemplated herein that a multi-compartment capsule of the present invention may comprise component parts of the capsule having various time-release coatings to facilitate the release and ultimately the absorption of those active ingredients introduced  
15 into the different receiving chambers of the multi-compartment capsule to release at different release rates. In particular, a primary capsule may be formed having a conventional time-release coating that dissolves and releases the active ingredient(s) contained therein before the timed-release of the active ingredient(s) contained within a secondary capsule. Likewise, the dividing walls disposed within the internal periphery of the base of a capsule  
20 may be formed having conventional time-release coatings that dissolve and release the active ingredients within each receiving chamber defined by the dividing walls at different rates, thereby delivering the active ingredients or medicaments contained within a multi-compartment capsule at different rates. Certain active ingredients or medicaments may, therefore, be delivered at a selected interval, while other ingredients may be released at  
25 a later interval. In this way, the novel design of the multi-compartment capsules of the present invention may facilitate precision delivery of active ingredients to targeted areas of the consumer.

The disclosure of secondary and tertiary capsules may be replaced with other forms of microencapsulation. Microencapsulation, as previously described, refers to the process whereby minute parcels of a solid, liquid, gas or dispersion, introduced into one or more of the receiving chambers as active ingredient(s), are film-coated with a secondary material in order to shield the active ingredient from its surrounding environment. Microcapsules may measure from microns to several millimeters, whereas the main purpose being to facilitate the release of the active ingredients at different release rates.

The incorporation of time-release coatings to varying the release rates of the active ingredients of a multi-compartment capsule may be used to target key time intervals or events when the body may be most able to utilize the active ingredients. In one presently preferred embodiment of the present invention, all of the active ingredients may be microencapsulated. In alternate presently preferred embodiments, only selected ingredients may be microencapsulated for delayed release, while other ingredients may be provided for immediate absorption. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

As contemplated herein, the physical states of active ingredients are characterized into one of four different states (*e.g.*, solid, liquid, gas or dispersion). These four different states are sometimes referred to as "phases" (*i.e.*, solid phase, liquid phase, gas phase or dispersion phase). For purposes of the present invention, the term "solid" is defined as including, by way of example only and not by way of limitation, pills, tablets, capsules (including both hard and soft elastic), powders, granulation, flakes, troches (lozenges and pastilles), suppositories and semi-solid pastes, ointments, emulsions or creams. The term "liquid" is defined as including, by way of example only and not by way of limitation, solutions, spirits, elixirs, sprays, syrups or fluid extracts. The term "dispersion" is defined as including, by way of example only and not by way of limitation, aerosols (liquid or solid in gas), suspensions



(solid in liquid), emulsions (liquid in liquid), foams (gas in liquid), solid foams (solid in gas) or gels (liquid or solid in solid).

The active ingredients or medicaments introduced into the receiving chambers of the multi-compartment capsules of the present invention preferably comprise a pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof. For purposes of the present invention, the term "pharmaceutical" is defined as any substance that affects the structure or functioning of a living organism. Pharmaceuticals, sometimes referred to as "drugs" are widely used for the prevention, diagnosis and treatment of diseases and for the relief of symptoms. The term "biotechnical" is defined as any substance that is derived from a biotechnology process. Biotechnology, sometimes shortened to "biotech", is the development of techniques and methods (*e.g.*, genetic engineering, protein engineering, genomics, proteomics, monoclonal antibody production, polymerase chain reaction, transgenics and the like) for the application of biological processes to the production of materials of use in medicine, foods, nutrition and industry. The term "nutraceutical" is defined as any substance that is a food or a part of a food and provides medical or health benefits, including the prevention and treatment of disease. The term "vitamin" is defined as any of various organic substances or compounds that are essential for the normal processes of growth and maintenance (*e.g.*, essential for energy transformation and regulation of metabolism) of the body which are present in natural foodstuffs or sometimes produced within the body. The term "dietary supplement" is defined as any product (other than tobacco) intended to supplement the diet that bears or contains one or more of the following dietary ingredients: (A) a vitamin; (B) a mineral; (C) an herb or other botanical; (D) an amino acid; (E) a dietary substance for supplementing the diet by increasing the total dietary intake; or (F) a concentrate, metabolite, constituent, extract or combination of any ingredient described in (A), (B), (C), (D), or (E) hereinabove. If desired, excipients may also be introduced into one or more of the receiving chambers of the multi-compartment capsules of the present invention in addition to the active ingredient(s). For example, in some cases involving medicaments with poor water solubility, it may be desirable to stabilize the liquids,

solids or dispersions using a lipid, lipoid, lecithin, ghee or the like. Still further by example, in some cases involving active ingredients or medicaments with poor bioavailability, bioequivalence, or other undesirous pharmaceutical properties (*e.g.*, poor water solubility, pH lability, physical incompatibility, chemical incompatibility, macromolecular size and the like) such as proteins (*e.g.*, hormones, erythropoietins, colony stimulating factors, interferons, interleukins, plasminogen activators, monoclonal antibodies, vaccines, plant proteins, such as soy and other therapeutic proteins) or other non-polar or weak-polar materials, it may be desirable to complement the active ingredient or medicament in liquid, solid or dispersion form using a fat, lipid, lipoid, lecithin, ghee, polymers, viral and bacterial vectors and the like.

\* \* \* \* \*

It may be demonstrated that as medical and pharmacy knowledge has continued to expand exponentially, new medicaments, new classes of medicaments and new delivery technologies are becoming available for use in animals and humans who experience particular medical diseases, illnesses or conditions. A disease, illness, or condition may affect one or more organ systems in an animal or human. Organ systems may include, for example: (1) autonomic, (2) cardiovascular, (3) neurological, (4) gastro-intestinal, (5) respiratory, (6) renal system, (7) psychiatric, (8) endocrine, (9) gynecologic, (10) urologic, (11) immunologic, (12) bone and joint systems, (13) ear, nose, and throat, (14) dermatologic, (15) hematologic, (16) infectious defense and (17) nutrition and metabolism. In an animal or human who may be suffering from one disease, illness or condition, it is common to also be suffering from a disease, illness or condition affecting one or more of the other organ system(s). These concomitant diseases, illnesses or conditions occurring within a single animal or human are often labeled as “co-morbidities,” a term often shortened and referred to as “co-morbid.”

New medicaments and delivery technologies are providing patients and their health care practitioners with unprecedented therapeutic options in managing diseases, illnesses and conditions. In spite of this sophistication, there has been no effort to develop new methods of

using fixed combinations of medicaments for therapy of co-morbid diseases, illnesses or conditions. Moreover, there has been no effort to develop new methods of using fixed combinations of medicaments for management of a single disease, illness or condition affecting one or more organ system(s). The aforementioned fixed combinations may include a plurality of medicaments, which may be newly discovered and developed, or have been known for sometime or some combination of medicaments thereof. In any regard, said fixed combinations have not previously been contemplated in the art.

\* \* \* \* \*

The following examples will illustrate the invention in further detail. It will be readily understood that the various active ingredients or medicaments that may be introduced into the receiving chambers of the multi-compartment capsules of the present invention, as generally described and illustrated in the Examples herein, are to be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or process for implementing those principles. Thus, the following more detailed description of the presently preferred embodiments of the methods, formulations, and compositions of the present invention, as represented in Examples I-XIV below, is not intended to limit the scope of the invention, as claimed, but is merely representative of presently preferred embodiments of the invention.

#### EXAMPLE I

[Glucosamine/Chondroitin (solid) & Vitamin E (liquid)]

As appreciated by those skilled in the art, arthritis is an inflammatory condition typically affecting the synovia membranes and cartilage of joints. It has been estimated that as many as one in three persons may experience symptoms associated with arthritis during their lifetime.

In addition to arthritis, various other chronic, debilitating conditions may afflict the aged. Many of these conditions result from the natural process of aging in humans. The natural aging process is partially due to the accumulation and effects of toxic free-radical

chemicals. Free-radicals result from several homeostatic biochemical processes. It is, accordingly, desirable to develop nutraceutical or dietary supplement products which may alleviate multiple chronic, debilitating conditions. It is also desirable to package and administer such products in the most economic and convenient possible fashion.

5           The administration of glucosamine, a naturally occurring substance in mucopoly-saccharides, mucoproteins and chitin, is believed to promote the production of cartilage components and the repair of damaged cartilage. Clinical findings support that fibroblast cells increased production of mucopolycaccharide and collagen synthesis when glucosamine was added.

10           Chondroitin sulfates are large polymers of glycosaminoglycans, primarily D-glucuronic acid and D-acetylgalactosamine, and disaccharides and may be derived from the cartilage of bovine trachea. The administration of chondroitin sulfate has been shown to promote the formation of new cartilage matrix. In particular, chondroitin stimulates the metabolism of chondrocyte cells and the production of collagen and proteoglycan.

15           Vitamin E, also known as alpha-tocopherol, is a well-known scavenger of free-radicals in the body. Free-radical scavengers are sometimes referred to as anti-oxidants. This scavenging process is important for detoxifying the body of chemicals which are known to promote apoptosis, or programmed cell death. Apoptosis is a scientific description of cellular destruction. Although vitamin E is a popular anti-oxidant, it is poorly soluble in  
20           water and thus can be administered only as a liquid-oil formulation or in an oil formulation enclosed in a soft elastic capsule.

          In one presently preferred embodiment of the present invention, therapeutically effective amounts of glucosamine, chondroitin, and vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein at least two of  
25           the active ingredients have physical states (*e.g.*, solid, liquid, gas or dispersion) that differ. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-arthritis

and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Primary Capsule:

5	Glucosamine HCl	500 mg
	[500-2000 mg/day]	
	Chondroitin sulfate	400 mg
	[400-1600 mg/day ]	

Secondary Capsule:

10	Vitamin E	200 IU
	[200 - 400 IU/day]	

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, glucosamine HCl/chondroitin sulfate and vitamin E) in the primary and secondary capsules, respectively, of the multi-compartment capsule may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

A therapeutically effective amount of glucosamine HCl/chondroitin sulfate may be introduced into at least a portion of the internal periphery of the receiving chambers of a primary capsule in the form of a solid and a therapeutically effective amount of vitamin E may be introduced into at least a portion of a secondary capsule in the form of a liquid, if desired. Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules,

respectively, is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. For example, a multi-compartment capsule may be formulated having glucosamine HCl and chondroitin sulfate introduced into the receiving chambers of the secondary capsule and vitamin E may be introduced into the receiving chamber of the primary capsule. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

## EXAMPLE II

[S-adenosylmethione (SAME) (solid) & Vitamin E (liquid)]

S-adenosylmethione (SAME), may be derived from two materials: methionine, a sulfur-containing amino acid, and adenosine triphosphate (ATP), the body's main energy compound. SAME was originally developed around 1950 as an antidepressant. Over the years, it has also been found that SAME may assist in alleviating arthritic symptoms, assist in the manufacture of melatonin, which is needed to regulate sleep, help protect DNA from harmful mutations and prevent certain types of nerve damage.

As noted above, vitamin E is a popular anti-oxidant, but it is poorly soluble in water and therefore can be administered only as a liquid-oil formulation. Vitamin E is typically measured in international units (IU) of alpha tocopherol.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of SAME and vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein SAME comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of vitamin E. As shown in Figures 3 and 4, a therapeutically effective amount of SAME may be introduced into receiving chamber 218a and a therapeutically effective amount of vitamin E may be introduced into receiving chamber 218b of a multi-compartment capsule 210 of the present invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for



effective amount of vitamin E, in the form of a liquid, may be introduced into receiving chamber 128 of a multi-compartment capsule 110 of the present invention. The material forming the primary capsule shell 111 may be formulated in a manner allowing for immediate dissolution and release of the contents of receiving chamber 118. The material forming the secondary capsule shell 120 may be formulated in a manner allowing for either an immediate dissolution or a time-delayed dissolution and release of the contents of receiving chamber 128. The material forming the tertiary capsule shell 138 may be formulated in a manner allowing for time-delayed dissolution and release of the contents of receiving chamber 138. In this presently preferred embodiment of the present invention, a total daily dosage of SAME may be delivered as two separate dosages within a single oral dosage form. One presently preferred embodiment of the present invention thus makes for a more convenient dosage form.

Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into receiving chambers defined within a capsule is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

### EXAMPLE III

[Curcumin, Holy Basil, Zinc (solid) & Fish Oil  
(Omega 3 Fatty Acids DHA & EPA - liquid)]

Curcumin belongs to a class of compounds derived from the turmeric root and is a yellow-orange volatile oil. It is believed that curcumin has an inhibitory effect on carcinogenesis, which is the evolution of a normal cell into a cancerous cell. There is clinical evidence to suggest curcumin may help to prevent stomach, colon, oral, esophageal, breast and skin cancers. Additional studies have been conducted to show that curcumin may be



helpful in balancing cholesterol levels, protecting against ulcers by inhibition of gastric acid secretion and protection of gastric mucosal tissue, and anti-inflammatory actions. In one clinical study, curcumin was found to be as effective as non-steroidal anti-inflammatory drugs in the treatment of arthritis and post-operative pain.

5        The administration of Holy Basil (*Ocimum sanctum*) has been shown to have an effect on promoting peripherally mediated analgesic effects. This action allows a broad range of therapeutic effects, including, anti-inflammatory, hypoglycemia, analgesic, anti-ulcer and anti-septic properties.

10        As known, zinc is a mineral that occurs in animal and plant tissues and is an important co-factor for various enzyme reactions in the body, as well as being helpful for the reproduction system, and for the manufacture of body proteins. Zinc is also an antioxidant nutrient, similar to vitamin E. There is clinical data that suggests that zinc may be important to the prostate and other reproductive organs in the body, may help in the contractility of muscles, help stabilize blood, help maintain the body's alkaline balance and aid in the  
15        digestion and metabolism of phosphorus.

Over several decades considerable evidence has been collected to suggest that fish and fish oils are beneficial to the heart, mental health and in reducing cancer risk. The "active" components of fish oils are eicosapentaenoic acid (EPA), a polyunsaturated fatty acid with a 20 carbon chain, and docosahexaenoic acid (DHA), a polyunsaturated fatty acid  
20        with a 22 carbon chain. Both active components are members of the omega-3 group of essential fatty acids and are found exclusively in marine animals. The best sources for EPA and DHA may be fatty fish such as herring, sardines, salmon and fresh tuna.

The recommended daily intake of EPA plus DHA is between 650 to 1000 mg/day. Clinical trials have used anywhere from 1 g/day to 10 g/day, but little additional benefit has  
25        been observed at levels above 5 g/day of EPA and DHA combined. The onset of beneficial effects is variable. Effects on cholesterol may occur in just a few weeks, but it may take there (3) months or longer to see effects in degenerative diseases, such as arthritis.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of curcumin, Holy Basil, zinc and fish oil (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein curcumin, Holy Basil and zinc comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of the fish oil. As shown in Figure 2, a therapeutically effective amount of curcumin may be introduced into receiving chamber 138 of a tertiary capsule 130, a therapeutically effective amount of Holy Basil and zinc may be introduced into receiving chamber 128 of a secondary capsule and a therapeutically effective amount of fish oil may be introduced into receiving chamber 118 of a primary capsule 111 of a multi-compartment capsule 110 of the present invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-neoplastic, anti-inflammatory, analgesic and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Tertiary Capsule (130):

Curcumin	400 mg
[1200-1800 mg/day; 400 mg three times daily ]	

Secondary Capsule (120):

Holy Basil	2.5 gms
[2.5 grams fresh dried leaf powder/day]	
Zinc	15 mg
[4 - 15 mg/day]	

Primary Capsule (111):

Fish oil	1000 mg
(Omega 3 fatty acids - DHA & EPA)	
[650 - 1000 mg/day]	

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, curcumin, Holy Basil, Zinc and fish oil) in the primary, secondary and tertiary capsules 111, 120, 130 of one presently preferred embodiment of a multi-compartment capsule 110 may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

As contemplated herein, a therapeutically effective amount of curcumin may be introduced into at least a portion of the receiving chamber 138 of the tertiary capsule 130 in the form of a solid, a therapeutically effective amount of Holy Basil and zinc may be introduced into at least a portion of the receiving chamber 128 of the secondary capsule 120 in the form of a solid and a therapeutically effective amount of fish oil may be introduced into at least a portion of the primary capsule 111 in the form of a liquid. Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules, respectively, is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

#### EXAMPLE IV

[Vitamin C (solid) & Vitamin E (liquid)]

It is believed that vitamin C plays an important role as a component of enzymes involved in the synthesis of collagen and carnitine. A vital role of vitamin C, however, is believed to be that of the primary, water-soluble antioxidant in the human body. A daily



amount of vitamin E may be introduced into at least a portion of a second receiving chamber in the form of a liquid. Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules, respectively, is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

#### EXAMPLE V

[Selenium/Vitamin C (solid) & Vitamin E,/Beta-Carotene/  
Fish Oil (Omega 3 Fatty Acids DHA & EPA) (liquid)]

Selenium is an essential trace mineral in the human body and an important part of antioxidant enzymes that protect cells against the effects of free radicals that are produced during normal oxygen metabolism. As readily known in the art, the body has developed defenses, such as antioxidants, to assist in controlling levels of free radicals which can cause damage to cells and contribute to the development of some chronic diseases. It is also believed that Selenium is essential for normal functioning of the immune system and thyroid gland. The recommended dietary allowance for selenium is 55 mcg/day.

As noted above, it is believed that vitamin C plays an important role as a component of enzymes involved in the synthesis of collagen and carnitine and a vital role as a water-soluble antioxidant in the human body. Vitamin E is another important anti-oxidant.

Beta-carotene is a substance found in plants that the body converts into vitamin A. It is believed that beta-carotene acts as an antioxidant and an immune system booster. There is no RDA for beta-carotene. The most common beta-carotene supplement intake is about 25,000 IU (15 mg) per day, however supplementation with as much as 100,000 IU (60 mg) per day has been reported.

It has been suggested that fish and fish oils are beneficial to the heart, mental health and in reducing cancer risk. The recommended daily intake of EPA plus DHA (the active components of fish oil) is between 650 to 1000 mg/day. Clinical trials have used anywhere from 1 g/day to 10 g/day, but little additional benefit has been observed at levels above 5 g/day of EPA and DHA combined.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of selenium, vitamin C, beta-carotene, vitamin E and fish oil (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein selenium and vitamin C comprise a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of vitamin E, beta-carotene and fish oil (omega 3 fatty acids DHA & EPA). Specifically, a therapeutically effective amount of selenium and vitamin C may be introduced into one or more receiving chambers of a primary capsule and a therapeutically effective amount of vitamin E, beta-carotene and fish oil (omega 3 fatty acids DHA & EPA) may be introduced into one or more receiving chambers of a secondary capsule to form a multi-compartment capsule of the present invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Primary Capsule:

Selenium	50 mcg
[50 - 100 mcg/day]	

Vitamin C	500 mg
[60-1000 mg/day ]	

Secondary Capsule:

Beta-carotene	50 mg
[30 - 300 mg/day]	

Vitamin E	200 IU
[200-400 IU/day]	

Fish oil  
(Omega 3 fatty acids - DHA & EPA)

1000 mg

5 [650 - 1000 mg/day]

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, selenium, vitamin C , vitamin E, beta carotene and fish oil) in different receiving chambers of a multi-compartment capsule may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment and is contemplated herein.

15 A therapeutically effective amount of selenium and vitamin C may be introduced into one or more receiving chambers of a primary capsule in solid form and a therapeutically effective amount of vitamin E, beta carotene and fish oil may be introduced into one or more receiving chambers of a secondary capsule in the form of a liquid. Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules, respectively, is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

#### EXAMPLE VI

[Fluoxetine (solid), S-adenosylmethione (SAME) (solid) & Vitamin E (liquid)]

30 As appreciated by those skilled in the art, depression is a mental state characterized by excessive sadness. Depression is one of several forms of mood disorders. Activity in those

affected with depression may be agitated and restless or slow and retarded. Those affected may also show pessimistic or despairing behavior and may have disturbances in sleep, appetite and concentration. Depression is often a co-morbid condition with other chronic disease states involving the neurological system, cardiovascular system, respiratory system, endocrine system, musculoskeletal system, immune system, genitourinary system and the like. This list is should not be considered exclusive.

Administration of Fluoxetine is known by those of skill in the art to alleviate the signs and symptoms of depression. Fluoxetine belongs to a class of compounds which are given the functional name: selective serotonin re-uptake inhibitors (SSRI's). This class may include, for example: fluoxetine (PROZAC®), sertraline (ZOLOFT®), paroxetine (PAXIL®), fluvoxamine (LUVOX®), citalopram (CELEXA®) and escitalopram (LEXAPRO®). As appreciated, the foregoing list is provided herein as exemplary and should not be considered exclusive or exhaustive.

Fluoxetine is a bicyclic compound, similar in structure to phenylpropanolamine. Fluoxetine structure imparts a high selectivity for interaction with cells of the nervous system for the function of preventing the re-uptake of serotonin into pre-synaptic cell storage sites. This action leads to marked increases in synaptic concentration of serotonin and is facilitative of numerous physiological processes requiring serotonin neurotransmission. In the pharmaceutical field Fluoxetine is available as a hydrochloride salt (HCl).

S-adenosylmethione (SAME), is derived from two materials: methionine, a sulfur-containing amino acid, and adenosine triphosphate (ATP), the body's main energy compound. SAME was originally developed around 1950 as an antidepressant, but it was also found to be helpful in the alleviation of arthritic symptoms. SAME is essential for the manufacture of melatonin, which is needed to regulate sleep. It also helps to protect DNA from harmful mutations and may help prevent certain types of nerve damage. Current clinical research is beginning to confirm these antidepressant qualities of SAME.

Vitamin E, also named alpha-tocopherol, is a well-known scavenger of free-radicals in the body. Free-radical scavengers are sometimes referred to as anti-oxidants. This



scavenging process is important for detoxifying the body of chemicals which are known to promote apoptosis, or programmed cell death. Apoptosis is a scientific description of cellular destruction. Although it is a popular anti-oxidant, Vitamin E is poorly soluble in water and thus can be administered only as a liquid-oil formulation or in an oil formulation enclosed in a soft elastic capsule. Vitamin E is typically measured in international units (IU) of alpha tocopherol.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of Fluoxetine, SAMe and Vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Fluoxetine and SAMe comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of Vitamin E. As shown in Figures 3 and 4, a therapeutically effective amount of Fluoxetine and SAMe may be introduced into receiving chamber 218a and a therapeutically effective amount of Vitamin E may be introduced into receiving chamber 218b of a multi-compartment capsule 210 of the present invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering mood enhancing, anti-depressant and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Receiving Chamber (218a):

Fluoxetine	20 mg
[20 - 60 mg/day]	
S-adenosylmethione	1000 mg
[200-1600 mg/day]	

Receiving Chamber (218b):

Vitamin E	200 IU
[200-400 IU/day]	

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, Fluoxetine/SAMe and Vitamin E) in the primary and secondary capsules, respectively, of the multi-compartment capsule may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

According to one presently preferred embodiment of the present invention, a therapeutically effective amount of Fluoxetine and SAMe may be introduced into at least a portion of the receiving chamber 218a in the form of a solid and a therapeutically effective amount of Vitamin E may be introduced into at least a portion of the receiving chamber 218b of the primary capsule 211 in the form of a liquid.

In an alternative presently preferred embodiment of the present invention, therapeutically effective amounts of Fluoxetine and SAMe and Vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Fluoxetine and SAMe comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of Vitamin E. As shown in Figure 2, a therapeutically effective amount of Fluoxetine and SAMe, in the form of a solid, may be introduced into receiving chamber 118 and 138 and a therapeutically effective amount of Vitamin E, in the form of a liquid, may be introduced into receiving chamber 128 of a multi-compartment capsule 110 of the present invention. The material forming the primary capsule shell 111 may be formulated in a manner allowing for immediate dissolution and release of the contents of receiving chamber 118. The material forming the secondary capsule shell 120 may be formulated in a manner allowing for either an immediate dissolution or a time-delayed dissolution and release of the contents of receiving chamber 128. The material forming the tertiary capsule shell 138 may be formulated in a manner allowing for time-delayed dissolution and release of the contents of receiving chamber 138. In this

presently preferred embodiment of the present invention, a total daily dosage of Fluoxetine and SAME may be delivered as two separate dosages within a single oral dosage form. One presently preferred embodiment of the present invention thus makes for a more convenient dosage form.

5            Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules, respectively, is anticipated such that the various ingredients may be introduced in different  
10   receiving chambers to accommodate different treatment modalities. For example, a multi-compartment capsule may be formulated having Fluoxetine and SAME introduced into the receiving chambers of the secondary capsule and Vitamin E may be introduced into the receiving chamber of the primary capsule. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as  
15   restrictive to a particular structure or method for implementing those principles.

#### EXAMPLE VII

##### [Rofecoxib (solid) & Vitamin E (liquid)]

As appreciated by those skilled in the art, arthritis is an inflammatory condition typically affecting the synovia and cartilage of joints. It has been estimated that as many as  
20   one in three persons may experience symptoms associated with arthritis during their lifetime.

In addition to arthritis, various other chronic, debilitating conditions may afflict the aged. Many of these conditions result from the natural process of aging in humans. The natural aging process is partially due to the accumulation and effects of toxic free-radical chemicals. Free-radicals result from several homeostatic biochemical processes. It is,  
25   accordingly, desirable to develop pharmaceutical, biotechnical, nutraceutical or dietary supplement products which may alleviate multiple chronic, debilitating conditions. It is also desirable to package and administer such products in the most economic and convenient possible fashion.

Anti-inflammatory agents may have many diverse therapeutic roles in the human body. Inflammation is the process undertaken by the body as it responds to an injury. A typical inflammatory response involves blood vessel dilation, increased blood flow to the site of injury, and influx of white blood cells to process and remove dead tissue. Inflammation can lead to pain and swelling at the site of injury. Medicaments used in modulating the inflammatory response may be divided into steroid and non-steroidal labels. The latter is more commonly identified as non-steroidal anti-inflammatory drugs (NSAIDs).

Rofecoxib belongs to a class of NSAID compounds given the functional name cyclo-oxygenase-2 ("COX-2") inhibitors. This class may include, for example: rofecoxib (VIOXX®), celecoxib (CELEBREX®), valdecoxib (BEXTRA®), and meloxicam (MOBIC®). As appreciated, the foregoing list is provided herein as exemplary and should not be considered exclusive or exhaustive.

Rofecoxib is presently believed to inhibit the action of COX-2, an enzyme involved in the production of prostaglandins in the human body. Prostaglandins serve many diverse roles, one of which is to stimulate an inflammation mechanism in immune responses. Recently, Rofecoxib was labeled for use in the treatment of osteoarthritis, rheumatoid arthritis, acute pain, and primary dysmenorrhea.

Vitamin E, also named alpha-tocopherol, is a well-known scavenger of free-radicals in the body. Free-radical scavengers are sometimes referred to as anti-oxidants. This scavenging process is important for detoxifying the body of chemicals which are known to promote apoptosis, or programmed cell death. Apoptosis is a scientific description of cellular destruction. Although it is a popular anti-oxidant, Vitamin E is poorly soluble in water and thus can be administered only as a liquid-oil formulation or in an oil formulation enclosed in a soft elastic capsule.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of Rofecoxib and Vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Rofecoxib comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of Vitamin E. As

shown in Figures 3 and 4, a therapeutically effective amount of Rofecoxib may be introduced into receiving chamber 218a and a therapeutically effective amount of Vitamin E may be introduced into receiving chamber 218b of a multi-compartment capsule 210 of the present invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-inflammatory and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Receiving Chamber (218a):

10	Rofecoxib	25 mg
	[12.5 - 25 mg/day]	

Receiving Chamber (218b):

	Vitamin E	200 IU
	[200-400 IU/day]	

15

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, Rofecoxib and Vitamin E) of the multi-compartment capsule 210 may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

20

According to one presently preferred embodiment of the present invention, a therapeutically effective amount of Rofecoxib may be introduced into at least a portion of the receiving chamber 218a in the form of a solid and a therapeutically effective amount of Vitamin E may be introduced into at least a portion of the receiving chamber 218b of the primary capsule 211 in the form of a liquid.

25

In an alternative presently preferred embodiment of the present invention, therapeutically effective amounts of Rofecoxib and Vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Rofecoxib comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of Vitamin E. As shown in Figure 2, a therapeutically effective amount of Rofecoxib, in the form of a solid, may be introduced into receiving chamber 118 and 138 and a therapeutically effective amount of Vitamin E, in the form of a liquid, may be introduced into receiving chamber 128 of a multi-compartment capsule 110 of the present invention. The material forming the primary capsule shell 111 may be formulated in a manner allowing for immediate dissolution and release of the contents of receiving chamber 118. The material forming the secondary capsule shell 120 may be formulated in a manner allowing for either an immediate dissolution or a time-delayed dissolution and release of the contents of receiving chamber 128. The material forming the tertiary capsule shell 138 may be formulated in a manner allowing for time-delayed dissolution and release of the contents of receiving chamber 138. In this presently preferred embodiment of the present invention, a total daily dosage of Rofecoxib may be delivered as two separate dosages within a single oral dosage form. One presently preferred embodiment of the present invention thus makes for a more convenient dosage form.

Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into receiving chambers defined within a capsule is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

### EXAMPLE VIII

[Diphenhydramine Hydrochloride (solid) & Vitamin E (liquid)]

As appreciated by those skilled in the art, allergic reactions are conditions wherein the immune system is stimulated to identify, segregate and dispose of exogenous chemicals which cannot be recognized by the body. Allergic reactions are often associated with the release of histamine, a chemical compound which produces changes in the permeability of blood vessels and the accumulation of other immune system cells. In some circumstances, it may be desirable to modulate the amount of allergic response that is capable of being generated by the immune system.

Diphenhydramine belongs to a class of compounds which are given the functional name: histamine-1 ( $H_1$ ) receptor antagonists. These compounds are more generally labeled as antihistamines. These antagonists are further divided according to their chemical structures. Diphenhydramine is an ethanolamine (aminoalkyl ether) derivative. Other chemical divisions may include, for example: ethylenediamine, propylamine, phenothiazine, piperazine. The ethanolamine division may include, for example: diphenhydramine, clemastine, dimenhydrinate, and doxylamine. As appreciated, the foregoing list is provided herein as exemplary and should not be considered exclusive or exhaustive.

Antihistamines block the interaction of the neurotransmitter, histamine, with  $H_1$  receptors located in smooth muscle linings of the gastrointestinal tract, bronchial tract and large blood vessels. This blocking action may lead to marked relaxation in smooth muscle tone and is facilitative of numerous physiological processes including respiration.

The  $H_1$  antagonists may also be divided according to their selectivity for central and peripheral  $H_1$  receptors. A second-generation of  $H_1$  has emerged in recent years. These agents have a greater selectivity for peripheral  $H_1$  receptors. Second-generation  $H_1$  receptor antagonists may include, for example: azelastine (ASTELIN<sup>®</sup>), cetirizine (Zyrtec<sup>®</sup>), desloratadine (Clarinet<sup>®</sup>), fexofenadine (Allegra<sup>®</sup>) and loratadine (Claritin<sup>®</sup>).

Vitamin E, also named alpha-tocopherol, is a well-known scavenger of free-radicals in the body. Free-radical scavengers are sometimes referred to as anti-oxidants. This

scavenging process is important for detoxifying the body of chemicals which are known to promote apoptosis, or programmed cell death. Apoptosis is a scientific description of cellular destruction. Although it is a popular anti-oxidant, Vitamin E is poorly soluble in water and thus can be administered only as a liquid-oil formulation or in an oil formulation enclosed in a soft elastic capsule.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of Diphenhydramine and Vitamin E (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein at least two of the active ingredients have physical states (*e.g.*, solid, liquid, gas or dispersion) that differ. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-allergic and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

Primary Capsule:

Diphenhydramine HCl	50 mg
[25 - 100 mg/day]	

Secondary Capsule:

Vitamin E	200 IU
[200-400 IU/day]	

The incorporation of time-release coatings to varying the release rates of the active ingredients ( *e.g.*, Diphenhydramine and Vitamin E) in the primary and secondary capsules, respectively, of the multi-compartment capsule may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.



A therapeutically effective amount of Diphenhydramine may be introduced into at least a portion of the internal periphery of the receiving chambers of a primary capsule in the form of a solid and a therapeutically effective amount of Vitamin E may be introduced into at least a portion of a secondary capsule in the form of a liquid, if desired. Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into the receiving chambers of the primary and secondary capsules, respectively, is anticipated such that the various ingredients may be introduced in different receiving chambers to accommodate different treatment modalities. For example, a multi-compartment capsule may be formulated having Diphenhydramine introduced into the receiving chambers of the secondary capsule and Vitamin E may be introduced into the receiving chamber of the primary capsule. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

#### EXAMPLE IX

[Celecoxib (solid) & Ibuprofen (liquid)]

As appreciated by those skilled in the art, arthritis is an inflammatory condition typically affecting the synovia and cartilage of joints. It has been estimated that as many as one in three persons may experience symptoms associated with arthritis during their lifetime.

Anti-inflammatory agents may have many diverse therapeutic roles in the human body. Inflammation is the process undertaken by the body as it responds to an injury. A typical inflammatory response involves blood vessel dilation, increased blood flow to the site of injury, and influx of white blood cells to process and remove dead tissue. Inflammation can lead to pain and swelling at the site of injury. Medicaments used in modulating the inflammatory response may be divided into steroid and non-steroidal labels. The latter is more commonly identified as non-steroidal anti-inflammatory drugs (NSAIDs).

Celecoxib belongs to a class of NSAID compounds given the functional name cyclo-oxygenase-2 ("COX-2") inhibitors. This class may include, for example: rofecoxib (VIOXX®), celecoxib (CELEBREX®), valdecoxib (BEXTRA®), etodolac (LODINE®) and meloxicam (MOBIC®). As appreciated, the foregoing list is provided herein as exemplary and should not be considered exclusive or exhaustive.

Celecoxib is believed to inhibit the action of COX-2, an enzyme involved in the production of prostaglandins in the human body. Prostaglandins serve many diverse roles, one of which is to stimulate an inflammation mechanism in immune responses. Recently, Celecoxib was labeled by the United States Food and Drug Administration (FDA) for use in the treatment of osteoarthritis, rheumatoid arthritis, acute pain, and primary dysmenorrhea.

Ibuprofen is another NSAID and is believed to function as a non-selective inhibitor of cyclo-oxygenase. Ibuprofen has been labeled by the FDA for use in the treatment of osteoarthritis, rheumatoid arthritis, relief of mild to moderate pain and primary dysmenorrhea. Ibuprofen belongs to a class of compounds called phenyl-a-methylacetic acids, which are derived from salicylic acid. Non-selective cyclo-oxygenase inhibitors may include, for example: ibuprofen (MOTRIN®), naproxen (NAPROSYN®), diclofenac (VOLTAREN®), flurbiprofen (ANSAID®), indomethacin (INDOCIN®), ketoprofen (ORUDIS®), ketorolac (TORADOL®), nabumetone (RELAFEN®), oxaprozin (DAYPRO®), piroxicam (FELDENE®) and sulindac (CLINORIL®). As appreciated, the foregoing list is provided herein as exemplary and should not be considered exclusive or exhaustive.

In one presently preferred embodiment of the present invention, therapeutically effective amounts of Celecoxib and Ibuprofen (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Celecoxib comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical state of Ibuprofen. As shown in Figures 3 and 4, a therapeutically effective amount of Celecoxib may be introduced into receiving chamber 218a and a therapeutically effective amount of Ibuprofen may be introduced into receiving chamber 218b of a multi-compartment capsule 210 of the present

invention. Consistent with the foregoing, multi-compartment, multi-phase capsules and encapsulation technology are herein contemplated to produce a delivery vehicle for delivering anti-arthritic and anti-oxidant compounds to the body in a single dosage. A capsular format of the present invention may include the following composition:

5

Receiving Chamber (218a):

Celecoxib	200 mg
[200 - 400 mg/day]	

Receiving Chamber (218b):

10

Ibuprofen	800 mg
[2400 - 3200 mg/day]	

15

The incorporation of time-release coatings to varying the release rates of the active ingredients (*e.g.*, Celecoxib and Ibuprofen) of the multi-compartment capsule 210 may be used to target key time intervals or events when the body may be most able to utilize the named active ingredients. Thus, the incorporation of time-release coatings in the encapsulation process when forming a multi-compartment capsule may be specifically designed to fit the needs and desires of numerous different users having similar conditions that are being targeted for treatment.

20

According to one presently preferred embodiment of the present invention, a therapeutically effective amount of Celecoxib may be introduced into at least a portion of the receiving chamber 218a in the form of a solid and a therapeutically effective amount of Ibuprofen may be introduced into at least a portion of the receiving chamber 218b of the primary capsule 211 in the form of a liquid.

25

In an alternative presently preferred embodiment of the present invention, therapeutically effective amounts of Celecoxib and Ibuprofen (active ingredients) may be introduced into receiving chambers of a multi-compartment capsule wherein Celecoxib comprises a physical state (*e.g.*, solid, liquid, gas or dispersion) different from the physical

state of Ibuprofen. As shown in Figure 2, a therapeutically effective amount of Celecoxib, in the form of a solid, may be introduced into receiving chamber 128 and a therapeutically effective amount of Ibuprofen, in the form of a liquid, may be introduced into receiving chambers 118 and 138 of a multi-compartment capsule 110 of the present invention.

5           The material forming the primary capsule shell 111 may be formulated in a manner allowing for immediate dissolution and release of the contents of receiving chamber 118. The material forming the secondary capsule shell 120 may be formulated in a manner allowing for either an immediate dissolution or a time-delayed dissolution and release of the contents of receiving chamber 128. The material forming the tertiary capsule shell 138 may  
10 be formulated in a manner allowing for time-delayed dissolution and release of the contents of receiving chamber 138. In this presently preferred embodiment of the present invention, a total daily dosage of Ibuprofen may be delivered as two separate dosages within a single oral dosage form. One presently preferred embodiment of the present invention thus makes for a more convenient dosage form.

15           Since the encapsulation process and multi-compartment, multi-phase capsule of the present invention are configured to apply to an anticipated treatment regime or medicinal design of a single dosage capsule, it will be readily appreciated that the introduction of one or more active ingredients into receiving chambers defined within a capsule is anticipated such that the various ingredients may be introduced in different receiving chambers to  
20 accommodate different treatment modalities. It is intended, therefore, that the examples provided herein be viewed as exemplary of the principles of the present invention, and not as restrictive to a particular structure or method for implementing those principles.

#### EXAMPLES X

25           Some embodiments of the present invention will use one or more of the below ingredients in a multi-compartment capsule, combinable as would be recognized in view of the teachings of the present application in combination with the knowledge available to one of ordinary skill in the art. It is noted that the following non-limiting lists illustrate

exemplary ingredeints that can be used with the present invention, including the the broader subclasses and classes to which they belong.

### **Botanicals**

Acerola Extracts	Alfalfa
Algae, Blue Green	Aloe
Amla	Angelica Root
Bacopa Monnieri	Mucuna Pruriens
Anise Seed	Arnica
Artichoke	Ashwagandha
Astragalus	Ayurvedic Herbs
Barberry	Barley Grass
Barley Sprout Extract	Benzoin
Bilberry	Bioflavonoids
Bitter Melon	Bitter Orange
Black Cohosh	Black Currant
Black Walnut	Bladderwrack
Blue Cohosh	Blueberry
Boswellia	Brahmi
Broccoli	Burdock
Butcher's Broom	Calendula
Capsicum	Cascara Sagrada
Cat's Claw	Cayenne
Celery Seed	Certified Organic Herbs
Chamomile	Chapparal
Chaste Berry	Chicory Root
Chinese Herbs	Chlorella
Chlorophyll	Citrus Aurantium
Cocoa	Coriander
Corn Silk	Cranberry
Curcuminoids	Damiana
Dandelion	Devil's Claw
Diosgenin	Dong Quai
Echinacea	Elderberry
Elecampane Root	Ephedra
Essential Oils	Eucalyptus
Evening Primrose	Eyebright
Fennel	Fenugreek
Feverfew	Flax Products
Fo Ti	Garcinia Cambogia
Garlic	Gentian
Ginger	Ginkgo Biloba
Ginseng (American)	Ginseng (Panax)
Ginseng (Siberian)	Goldenseal
Gotu Kola	Grape Seed Extract
Grape Skin Extract	Grapefruit Seed Extract
Green Food Products	Green Lipped Mussel Powder

Green Tea  
 Guarana  
 Gymnema Sylvestre  
 Henna  
 Herbal Teas  
 Horehound  
 Horsetail  
 Ipriflavone  
 Juniper Berries  
 Kelp Extract  
 Kombucha  
 Larch  
 Lemon Balm  
 Linden Flowers  
 Maca  
 Marshmallow  
 Molasses  
 Neem  
 Noni  
 Oatstraw  
 Olive Extract  
 Oregano Oil  
 Organic Sweeteners  
 Passion Flower  
 Pennyroyal  
 Pfaffia Paniculata  
 Piper Longum  
 Quercitin  
 Red Clover  
 Reishi Mushroom  
 Rhubarb Root  
 Rose Hips  
 Sage  
 Saw Palmetto  
 Seaweed extracts  
 Shatavari  
 Silymarin  
 Slippery Elm  
 Soybean Products  
 St. John's Wort  
 Summa  
 Terminalia Ajruna  
 Triphala  
 Uva Ursi  
 Vegetable Extracts  
 Wheat Germ  
 Wild Cherry bark  
 Witch Hazel  
 Yarrow  
 Yerba Sante

Griffonia Simplicifolia  
 Guggul  
 Hawthorne  
 Herbal Extracts, Standardized  
 Hops  
 Horse Chestnut  
 Hysop  
 Jojoba Oil  
 Kava Kava  
 Kola Nut  
 Kudzu  
 Lavender  
 Licorice Extract  
 Lobelia  
 Maitake Mushroom  
 Milk Thistle  
 Mushrooms  
 Nettle  
 Nopal  
 Octacosanol  
 Orange Peel Extract  
 Oregon Mountain Grape  
 Parsley  
 Pau d'Arco  
 Peppermint  
 Pine Bark Extract  
 Pygeum Africanum  
 Raspberry Powder  
 Red Raspberry  
 Resveratrol Extract  
 Rice Products  
 Rosemary Extract  
 Sarsaparilla  
 Schizandra  
 Senna  
 Shiitake Mushroom  
 Skullcap  
 Soy Isoflavones  
 Spirulina  
 Stevia  
 Tea Tree Oil  
 Tribulus Terrestris  
 Tumeric  
 Valerian Extract  
 Vitex  
 White Willow Bark  
 Wild Yam  
 Wormwood  
 Yellow Dock  
 Yohimbine

## Yucca

20-ECD 7-9%  
Acetyl L-Carnitine HCl 99%  
Alisma Extract 10:1  
Angelica Root Extract  
Artemisia Extract 4:1  
Asparagus Extract 4:1  
Astragalus Extract 10:1  
Astragalus Extract 5:1  
Astragalus Root Powder  
Avena Sativa Extract 10:1  
Barbed Skullcap Extract 10:1  
Bee Pollen Powder  
Bilberry Extract 10:1  
Black Cohosh Extract 2.5%  
Black Pepper Extract 4:1  
Bone Powder  
Broccoli Sprout Extract 10:1  
Buplerum (Chai Hu) Extract 5:1  
Cabbage Extract 4:1  
Caffeine 99%  
Calcium-Pyruvate 99%  
Cassia Nomame Extract 4:1  
Cat's Claw (Inner Bark) Powder  
Celandine (Greater) Extract 4:1  
Cetyl Myristoleate 11%  
Chaenomeles Extract 4:1  
Chamomile Flower Extract 4:1  
Chitin  
Chitosan 90%  
Chrysin 99%  
Cistanches Extract 5:1  
Citrus Bioflavonoid Complex 13%  
Clove Extract 5:1  
Coca Extract 4:1  
Colostrum  
Cordyceps Extract 7%  
Cornsilk Powder  
Cranberry Extract 4:1  
Curcumin Extract 95%  
Damiana Extract 4:1  
Dandelion Powder  
Danshen Extract 80%  
Devil's Claw Extract 2.5%  
Devil's Claw Root Powder  
Diosgenin 95%

## Extracts

4-Androstenedione 99%  
Adenophora Tetraohylla Ext 5:1  
Alpha Lipoic Acid 99%  
Arbutin 99%  
Artichoke Extract 5%, Globe  
Asparagus Powder  
Astragalus Extract 4:1  
Astragalus Root Extract 0.5%  
Atractylodes Extract 10:1  
Avena Sativa Extract 4:1  
Barberry Extract 10%  
Beta-Sisterol 35%  
Bitter Melon Extract 8:1  
Black Cohosh Root Powder  
Black Soy Bean Extract 10:1  
Boswellia Serrata Extract 65%  
Buchu Leaf Powder  
Burdock Root Extract 4:1  
Caffeine (Natural) 86-87%  
Calcium Citrate Granular 21%  
Carrot Root Extract 4:1  
Catnip Extract 4:1  
Cauliflower Extract 4:1  
Celery Seed Extract  
Cetyl Myristoleate 20%  
Chamomile Flower Extract 10:1  
ChasteTree Berry Extract 4:1  
Chitosan 80%  
Chondroitin Sulfate 90%  
Cinnamon Powder  
Citrus Aurantium Extract 6%  
Citrus Peel Extract 5:1  
Clove Powder  
Codonopsis Pilosula Extract 5:1  
Common Peony Extract 8:1  
Cornsilk Extract 4:1  
Corydalis Extract 10:1  
Cranberry Powder  
Cuscuta Extract 5:1  
Damiana Leaves Powder  
Dandelion Root Extract 6:1  
D-Calcium Pantothenate  
Devil's Claw Extract 4:1  
DHEA 99%  
DL-Phenyl Alanine

DMAE Bitartrate  
 Dong Quai Extract 4:1  
 D-Ribose  
 Echinacea Leaf Powder  
 Echinacea Purpurea Extract 4%  
 Echinacea Purpurea Root Powder  
 Elderberry Extract 20:1  
 Epimedium Extract 10%  
 Epimedium Extract 4:1  
 Epimedium Powder  
 Fennel Seed Extract 4:1  
 Fenugreek Extract 4:1  
 Feverfew Extract 5:1  
 Fish Oil Powder  
 Forskolin 8%  
 Fo-Ti Extract 8:1  
 Gardenia Extract 8:1  
 Garlic Powder  
 Ginger Extract 4:1  
 Ginger Root Powder  
 Ginkgo Extract 24/6%  
 Ginkgo Extract 24/7%  
 Ginkgo Leaf Powder  
 Ginseng (Panax) Extract 5%  
 Ginseng (Panax) Extract 80%  
 Glucosamine HCl 95% Granulation  
 Glucosamine Sulfate Potassium  
 Glucosamine Sulfate Sodium 99%  
 Goldenrod Powder  
 Goldenseal Root Powder  
 Gotu Kola Extract 4:1  
 Gotu Kola Powder  
 Grape Seed  
 Grape Seed Extract 20:1  
 Grape Seed Extract 5:1  
 Grape Seed Powder  
 Grape Skin Extract 4:1  
 Green Lip Mussel Extract  
 Green Tea Extract 4:1  
 Guarana Seed Extract 10%  
 Guarana Seed Extract 25%  
 Guggul Extract 2.5%  
 Gymnema Sylvestre Extract 25%  
 Hawthorne Berry Extract 4:1  
 Hawthorne Leaf Extract 2%  
 Hesperidin Extract 98%  
 Hops Flower Extract 4:1  
 Horehound Extract 4:1  
 Horse Chestnut Extract 20%  
 Horse Chestnut Powder

Dong Quai Extract 10:1  
 Dong Quai Root Powder  
 Echinacea Angustifolia Extract 4:1  
 Echinacea Purpurea Extract 10:1  
 Echinacea Purpurea Extract 4:1  
 Elder Flower Extract 4:1  
 Elderberry Extract 4:1  
 Epimedium Extract 10:1  
 Epimedium Extract 5%  
 Eucommia (Du Zhong) Extract 5:1  
 Fennel Seed Powder  
 Fenugreek Extract 6:1  
 Fisetin  
 Forbidden Palace Flower Extract 5:1  
 Fo-Ti Extract 12:1  
 Fo-Ti Powder  
 Garlic Extract 4:1  
 Gentian Root Extract 6:1  
 Ginger Root Extract 5%  
 Ginkgo Biloba Extract 8:1  
 Ginkgo Extract 24/6%<5  
 Ginkgo Leaf Extract 4:1  
 Ginseng (Korean) Powder  
 Ginseng (Panax) Extract 8%  
 Glucomannans Konjac Powder  
 Glucosamine HCl 99%  
 Glucosamine Sulfate Sodium 95% Granulation  
 Goldenrod Extract 4:1  
 Goldenseal Root Extract 14%  
 Gotu Kola Extract 16%  
 Gotu Kola Extract 8:1  
 Grape Fruit Powder  
 Grape Seed Extract 10:1  
 Grape Seed Extract 4:1  
 Grape Seed Extract 95%  
 Grape Skin Extract 20:1  
 Grass-Leaved Sweetflai Extract  
 Green Tea Extract 30%  
 Green Tea Extract 95%  
 Guarana Seed Extract 22%  
 Guggul Extract 10%  
 Gugulipid Extract 10%  
 Gymnema Sylvestre Powder  
 Hawthorne Berry Powder  
 Hearbacious Peony Extract 5:1  
 Honeysuckle Herb Extract 4:1  
 Horehound Extract 10:1  
 Horehound Herb Powder  
 Horse Chestnut Extract 4:1  
 Horsetail Extract 7%



Horsetail Powder  
 Hydrangea Extract 8:1  
 Hyssop Extract 4:1  
 Isodon Glaucoalyx Extract 10:1  
 Jiaogulan Extract 4:1  
 Jingjie Extract 4:1  
 Kava Kava Extract 30%  
 Kelp Extract 4:1  
 Kidney Bean Extract 10:1  
 Kidney Bean Pole 8:1  
 Kola Nut Extract 10%  
 Kudzu Extract 6:1  
 L-Glutamine  
 Licorice Extract 10%  
 Licorice Powder  
 L-Tyrosine  
 Lycium Fruit Extract 5:1  
 Ma Huang Extract 8%  
 Maca Extract 4:1  
 Magnesium Stearate  
 Magnolia Officinal Extract 4:1  
 Marigold Extract (Lutein 5%)  
 Methylsulfonylmethane 99%  
 Milk Thistle Seed Extract 80% silymarin  
 Motherwort Extract 4:1  
 Mucuna Pruriens Extract (15% L-Dopa)  
 Muira Puama Extract 4:1  
 Mushroom Extract 10:1 (feishi)  
 Myrobalan Extract 4:1  
 N-Acetyl-D-Glucosamine  
 Nettle Extract 7%  
 Nettle Leaf Powder  
 Olive Leaf Extract 18%  
 Orange Peel Extract 4:1  
 Oroxylum Indicum Extract 4:1  
 Oyster Meat Powder  
 Papaya Fruit Extract 4:1  
 Parsley Extract 4:1  
 Parsley Powder  
 Passion Flower Powder  
 Peppermint Extract 4:1  
 Perilla Seed Extract 4:1  
 Pharbitidis Extract 4:1  
 Pine Bark Extract 4:1  
 Polygala Tenoifolia Extract 4:1  
 Polygonum Extract 4:1  
 Propolis Extract 3%  
 Psyllium extract 4:1  
 Purple Willow Bark Extract 4:1  
 Pygeum Extract 4:1

Houltuynia Cordata Extract 5:1  
 Hydroxy Apatite  
 Indole-3-Carbinol 99%  
 Japanese Knotweed Extract  
 Jin Qian Cao Extract 4:1  
 Jujube Fruits Extract 4:1  
 Kava Kava Powder  
 Kelp Powder  
 Kidney Bean Pole 4:1  
 Kidney Bean Powder  
 Kudzu Extract 4:1  
 Lettuce Extract 4:1  
 L-Glycine  
 Licorice Extract 5:1  
 Lotus Leaf Powder  
 Lycium Fruit Extract 4:1  
 Ma Huang Extract 6%  
 Maca Extract 0.6%  
 Maca Root Powder  
 Magnolia Bark Powder  
 Maitake Mushroom Extract 4:1  
 Methoxyisoflavone 99%  
 Milk Thistle Extract 4:1  
 Morinda Extract 5:1  
 Motherwort Powder  
 Muira Puama Extract 12:1  
 Muira Puama Powder  
 Mustard Seed Extract 8:1  
 Myrrha Gum Extract 2.5%  
 N-Acetyl-L-Cysteine  
 Nettle Leaf Extract 4:1  
 Noni Powder  
 Olive Powder  
 Orange Peel Powder  
 Oroxylum Indicum Powder  
 Oyster Shell Powder  
 Parsley Extract 10:1  
 Parsley Leaf Extract 4:1  
 Passion Flower Extract 4:1  
 Pau D'Arco Powder  
 Peppermint Powder  
 Periwinkle Extract 4:1  
 Phosphatidyl Serine 20%  
 Plantago Asiatica Leaf Extract 5:1  
 Polygonum Extract  
 Pregnenolone 99%  
 Pseudoginseng Extract  
 Pumpkin Seed Extract 4:1  
 Purslane Herb Extract 4:1  
 Quercetin

Radish Extract 4:1  
 Radix Polygoni Extract 4:1  
 Red Pepper Extract 4:1  
 Red Yeast Rice Extract 10:1  
 Rehmannia Root Extract 4:1  
 Rhodiola Rosea Extract 4:1  
 Rhododendron Powder  
 Rhubarb Root Powder  
 Rice Powder  
 Rumex Madaid Extract 4:1  
 Salvia Extract 4:1  
 Saw Palmetto Extract 25%  
 Saw Palmetto Extract 25%  
 Saw Palmetto Extract 45-50%  
 Saw Palmetto Powder  
 Schizandra Extract 4:1  
 Sea Cucumber Powder  
 Sesame (Black) Seed Powder  
 Shitake Mushroom Extract  
 Siberian Ginseng Extract 4:1  
 Skullcap Extract 4:1  
 Slippery Elm Powder  
 Songaria Cynomorium Extract 4:1  
 Spirulina Powder  
 St. John's Wort Extract 4:1  
 Stanol 50%  
 Stevia Extract 4:1  
 Suma Root Extract 4:1  
 Taurine Powder  
 Tomato Extract  
 (trans)-Resveratrol 20-25%  
 Tribulus Extract 40%  
 Trifal Extract 4:1  
 Turmeric Root Powder  
 Uva Ursi Powder  
 Valerian Root Extract 4:1  
 Vinca Major Seed Extract 10:1  
 White Willow Bark 15% (total salicins)  
 White Willow Bark 25%  
 White Willow Bark Powder  
 Wild Yam Extract 16%  
 Wild Yam Extract 6%  
 Williams Elder Extract 4:1  
 Wolfiporia Extract 8:1  
 Yerba Mate Extract (2% caffeine)  
 Yohimbe Bark Extract 15:1  
 Yohimbe Bark Extract 3%  
 Yucca Extract 4:1

Radix Isatidis Extract 4:1  
 Red Clover Extract 4:1  
 Red Yeast Rice  
 Red Yeast Rice Powder  
 Reishi Mushroom Extract 4:1  
 Rhododendron Extract 4:1  
 Rhubarb Extract 4:1  
 Riboflavin (B2)  
 Rosemary Extract 20%  
 Salvia Extract 10:1  
 SAME  
 Saw Palmetto Extract 25%  
 Saw Palmetto Extract 4:1  
 Saw Palmetto Oil 85-95%  
 Schizandra Extract 10:1  
 Scopolia Acutangula Powder  
 Senna Leaf Powder  
 Shark Cartilage Powder  
 Siberian Ginseng Extract 0.8%  
 Siberian Ginseng Powder  
 Skullcap Extract 4:1  
 Sodium-Pyruvate 99%  
 Songaricum Powder  
 St. John's Wort Extract 0.3%  
 St. John's Wort Powder  
 Stephania Extract 4:1  
 Sulfate N+  
 Suma Root Powder  
 Thorowax Extract 4:1  
 Tomato Extract (0.2% Lycopene)  
 Tribulus Extract 10:1  
 Tribulus Powder  
 Turmeric Extract 4:1  
 Uva Ursi Extract 4:1  
 Valerian Root Extract 0.8%  
 Valerian Root Powder  
 White Wax Extract 4:1  
 White Willow Bark 20%  
 White Willow Bark Extract 4:1  
 Wild Yam Extract 10:1  
 Wild Yam Extract 4:1  
 Wild Yam Powder  
 Wolfberry Fruit Extract 10:1  
 Yellow Dock Root Extract 4:1  
 Yerba Mate Extract 4:1  
 Yohimbe Bark Extract 2%  
 Yohimbe Bark Powder

## Enzymes

Alpha Galactosidase  
Bromelain  
Papain  
Protease  
Superoxide Dismutase

Amylase  
Cellulase  
Peptidase  
Proteolytic Enzymes  
Trypsin

## Phospholipids

Lecithin  
Phosphatidyl Serine

Phosphatidyl Choline

5

## Specialty Nutraceuticals

5-Hydroxytryptophan  
Alpha Lipoic Acid  
Bee Products  
Bovine Cartilage  
Cetyl Myristoleate  
Chitosan  
Chondroitin Sulfate  
Collagen  
Creatine  
DMAE  
Germanium Sesquioxide  
Glucosamine HCL  
HMB (Hydroxyl Methyl Butyrate)  
Lactic Acid  
Liver Products  
Maltose-anhydrous  
MSM  
Phytosterols  
Pyruvate  
SAmE  
Shark Cartilage  
Vanadyl Sulfate  
Yeast

Acetyl L-Carnitine  
Alpha-Ketoglutarates  
Betaine Hydrochloride  
Caffeine  
Charcoal  
Choline  
Coenzyme Q10  
Colostrum  
Cyanocobalamin (Vitamin B12)  
Fumaric Acid  
Glandular Products  
Glucosamine Sulfate  
Immunoglobulin (Immune System Support)  
L-Carnitine  
Malic Acid  
Mannose (d-mannose)  
Other Carnitine Products  
Picolinic Acid  
Red Yeast Extract  
Selenium Yeast  
Theobromine  
Velvet Deer Antler

10

## Herbal Oils

Aloe Vera  
Artichoke Oil  
Black Currant Seed Oil 15% GLA  
Borage Oil 22% GLA

Artichoke Oil  
Black Currant Seed Oil 14% GLA  
Borage Oil 20% GLA  
Boswellia Serrata Oil

CLA Conjugated Linolic Acid 75% min.  
 Evening Primrose Oil 9% GLA  
 Garlic Oil  
 Guggul Lipid Oil  
 Oregano Oil  
 Pumpkin Seed Oil  
 Rosehip Oil  
 Saw Palmetto Oil  
 Tocotrienol Palm Oil  
 Wheat Germ Oil

Evening Primrose Oil 10% GLA  
 Flax Seed Oil 50% ALA  
 Grape Seed Oil  
 Olive Leaf Extract  
 Perilla Oil 60% ALA  
 Pygeum Oil  
 Rosemary Oil  
 Sterols  
 Walnut Oil  
 Sesame Seed Oil  
 Dill Seed Oil  
 Clove Bud Oil  
 Ginger Root Oil  
 Cinnamon Leaf Oil  
 Fennel Seed Oil  
 Curcuma Longa Oil  
 Cummin Seed Oil  
 Celery Seed Oil  
 Coriander Seed Oil  
  
 Red Raspberry Seed Oil  
 Cranberry Seed Oil  
 Blackberry Seed Oil

#### **Marine Oils**

Cod Liver Oil ( 1000 A /100 D )  
 Fish Oil 30% EPA / 20% DHA  
 Fish Oil Deodorized  
 Marine Lipid Oil 30/20  
 Salmon Oil 18% EPA / 12% DHA

Cod Liver Oil ( 2500A / 250D )  
 Fish Oil Concentrated  
 Marine Lipid Oil 18/12  
 Marine Lipid Oil 36/24  
 Squalene Oil ( Shark )

5

#### **Other Oils**

Alpha Lipoic Acid  
 Coenzyme Q10 .  
 Medium Chain Triglycerides MCT .

Cetyl Myristoleate CM  
 Lecithin

#### **Vitamins**

Ascorbic Acid (Vitamin C)  
 Biotin  
 Folic Acid  
 Inositol  
 Mixed Tocopherols  
 Orotic Acid  
 Pantothenates  
 Pyridoxine Hydrochloride (Vitamin B6)

B Vitamins  
 Fat Soluble Vitamins  
 HCA (Hydroxycitric Acid)  
 Mineral Ascorbates  
 Niacin (Vitamin B3)  
 PABA (Para-Aminobenzoic Acid)  
 Pantothenic Acid (Vitamin B5)  
 Riboflavin (Vitamin B2)

Synthetic Vitamins  
Tocotrienols  
Vitamin D  
Vitamin F  
Vitamin Oils  
Vitamin-Mineral Premixes

Thiamine (Vitamin B1)  
Vitamin A  
Vitamin E  
Vitamin K  
Vitamin Premixes  
Water Soluble Vitamins

### **Carotenoids**

Apocarotenal  
Beta-Carotene  
Carotenoids  
Lycopene

Astaxanthin  
Canthaxanthin  
Lutein / Lutein Esters  
Zeaxanthin

5

### **Hormones**

7-Keto-DHEA  
DHEA  
Nor-Androstenedione  
Progesterone

Androstenedione  
Melatonin  
Pregnenolone  
19 Nor-4-Androstendiol  
19 Nor-4-Androstenedione  
19 Nor-5-Androstenediol  
19 Nor-5-Androstendione  
3-Indolebutyric Acid  
4 Androstendiol  
4 Androstendione  
6 Furfurylaminopurine  
6-Benzylaminopurine

### **Minerals**

Boron  
Chelated Minerals  
Chromium  
Cobalt  
Dolomite  
Iron  
Manganese  
Mineral Products  
Other Minerals  
Potassium  
Sodium  
Trace Minerals  
Zinc

Calcium  
Chloride  
Coated Minerals  
Copper  
Iodine  
Magnesium  
Mineral Premixes  
Molybdenum  
Phosphorus  
Selenium  
Specialty Minerals  
Vanadium  
Malic Acid  
Pyruvate

10

## Probiotics

Acidophilus  
Lactobacillus

Bifido Bacteria

## Proteins / Amino Acids

Amino Acids  
Casein  
Glutamic Acid  
L-Arginine  
L-Glutamine  
L-Histidine  
L-Leucine  
L-Methionine  
L-Phenylalanine  
L-Taurine  
L-Tryptophan  
L-Valine  
Protein  
Soy Protein Isolates  
Whey Protein Isolates

Betaine  
Functional Soy  
L-Alanine  
L-Cysteine  
L-Glycine  
L-Isoeucince  
L-Lysine  
L-Ornithine  
L-Proline  
L-Threonine  
L-Tyrosine  
N-Acetyl-L-Cysteine  
Soluble Soy  
Textured Soy

5

## Specialty Nutrients

ATP  
Forskolin  
10 Sterol Esters  
Stanol Esters  
Probiotics  
Lactoferrin  
Lutein Esters  
15 Zeaxanthin  
Immunoglobulins  
Ipriflavone  
Isoflavones  
Fructo-Oligo-Saccharides  
20 Inulin  
Huperzine A  
Melatonin  
Medicinal Mushrooms  
Bile Products  
25 Peptone Products  
Glandular Products  
Pancreatic Products  
Thyroid Products



2-PAM: Acetylcholinesterase Reactivator Pralidoxime (Protopam) {2-PAM}: peripheral acetylcholinesterase reactivator for certain phosphoryl-enzyme complexes

Ganglionic Blockers

- 5
  - hexamethonium
  - mecamylamine (Inversine)
  - trimethaphan

Catecholamines

- 10
  - dobutamine (Dobutrex)
  - dopamine (Intropin)
  - epinephrine
  - isoproterenol (Isuprel)
  - norepinephrine (Levophed)

Direct Adrenoceptor Agonist Medicaments

- 15
  - albuterol (Ventolin, Provertil)
  - clonidine (Catapres)
  - methoxamine (Vasoxyl)
  - oxymetazoline (Afrin)
  - phenylephrine (Neo-Synephrine)
- 20
  - ritodrine (Yutopar)
  - salmeterol (Serevent)
  - terbutaline (Brethine)

Indirect-Acting Sympathomimetic Medicaments

- 25
  - amphetamine
  - cocaine
  - ephedrine, Pseudoephedrine tyramine

Alpha-Adrenoceptor Antagonists Medicaments

- 30
  - doxazosin (Cardura)
  - labetalol (Trandate, Normodyne)
  - phenoxybenzamine (Dibenzylamine)
  - phentolamine (Regitine)
  - prazosin (Minipress)
  - terazosin (Hytrin)
  - tolazoline (Priscoline)
- 35
  - trimazosin
  - yohimbine (Yocon)

$\beta$ -Adrenoceptor antagonist Medicaments

- 40
  - atenolol (Tenormin)
  - butoxamine
  - esmolol (Brevibloc)
  - labetalol (Trandate, Normodyne)
  - metoprolol (Lopressor)
  - nadolol (Corgard)
  - pindolol (Visken)
- 45
  - propranolol (Inderal)
  - timolol (Blocadren)

Adrenergic Neuron Blocking Medicaments

- 50
  - guanethidine (Ismelin)
  - reserpine



## **CARDIOVASCULAR SYSTEM DISORDERS**

	Cardiovascular testing and diagnosis
	Hypertension (HTN)
	Heart Failure
5	Ischemic Heart Disease
	Myocardial Infarction
	Arrhythmias
	Isolated Diastolic Heart Failure and Cardiomyopathies
	Cardiac Transplantation
10	Venous Thromboembolism
	Stroke
	Hyperlipidemia
	Peripheral vascular disease
15	Diuretics
	carbonic-anhydrase inhibitors
	loop diuretics
	osmotic diuretics
	potassium sparing diuretics
20	thiazide diuretics
	<b>Anitiarrhythmic Medicaments</b>
	Sodium Channel blocking agents
25	isopyramide (Norpace)
	flecainide (Tambocor)
	ibutilide
	lidocaine (Xylocaine)
	mexiletine (Mexitol)
	moricizine (Ethmozine)
30	procainamide (Pronestyl, Procan)
	propafenone (Rythmol)
	quinidine
	tocainide (Tonocard)
	Calcium Channel blocking agents
35	bepridil (Vasocor)
	diltiazem (Cardizem)
	verapamil (Isoptin, Calan)
	Adrenergic receptor antagonists
	propranolol (Inderal)
40	Other medicaments
	adenosine (Adenocard)
	amiodarone (Cordarone)
	bretylum (Bretylol)
	disopyramide (Norpace)
45	esmolol (Brevibloc)
	sotalol (Betapace)
	<b>Hypolipidemic medicaments</b>
	HMG CoA Reductase Inhibitors
	atorvastatin (Lipitor)
50	cerivistatin (Baycol)

	lovastatin (Mevacor)
	pravastatin (Pravochol)
	simvastatin (Zocor)
	Bile-acid sequestrants
5	cholestyramine (Questran)
	colestipol (Colestid)
	Fibric acids
	clofilbrate
	fenofibrate (Tricor)
10	gemfibrozil (Lopid)
	niacin, nicotinic acid
	probucol (Lorelco)
	<b>Antihypertensive medicaments</b>
	Adrenergic receptor antagonists
15	acebutalol (Sectral)
	atenolol (Tenormin)
	betaxolol (Betoptic)
	bisoprolol (Zebeta)
	carteolol (Cartrol)
20	clonidine (Catapres)
	labetalol (Normodyne)
	metoprolol (Toprol)
	penbutalol (Levatol)
	pindolol (Visken)
25	prazosin (Minipres)
	propranolol (Inderal)
	terazosin (Hytrin)
	timolol (Timoptic)
	Calcium Channel Antagonists
30	amlodipine (Norvasc)
	diltiazem (Cardizem)
	felodipine (Plendil)
	isradipine (Dynacirc)
	nicardipine (Cardene)
35	nifedipine (Procardia)
	nimodipine (Nimotop)
	nisoldipine (Sular)
	verapamil (Isoptin, Catan)
	Angiotensin Converting Enzyme (ACE) Inhibitor
40	benazepril (Lotensin)
	bepridil (Vascor)
	captopril (Capoten)
	enalapril (Vasotec)
	fosinopril (Monopril)
45	lisinopril (Prinivil, Zestril)
	moexipril (Univasc)
	quinapril (Accupril)
	ramipril (Altace)
	Angiotensin II Receptor Antagonists
50	losartan (Cozaar)

	valasartan (Diovan)
	<b>Diuretics</b>
5	amiloride (Midamor)
	bumetanide (Bumex)
	chlorothalidone (Hygroton)
	ethacrynic acid (Edecrin)
	furosemide (Lasix)
	hydrochlorothiazide (Diuril)
10	indapamide (Lozol)
	metolazone (Zaroxolyn)
	torsemide (Demadex)
	triamterene
	<b>Other Agents</b>
15	hydralazine (Apresoline)
	minoxidil (Rogaine)
	nitroprusside (Nipride)
	prazosin (Minipres)
	reserpine
20	sotalol (Brevibloc)
	spironolactone (Aldactone)
	terazosin (Hytrin)
	<b>Antianginal medicaments</b>
	Organic nitrates
25	Calcium Channel Antagonists
	Adrenergic Receptor Antagonists
	amyl nitrite
	erythrityl tetranitrate
	isosorbide dinitrate (Isordil)
30	nitroglycerin
	pentaerythritol tetranitrate
	<b>Congestive Heart Failure Medicaments</b>
	phosphodiesterase (PDE) inhibitors
	amrinone (Inocor)
35	milrinone (Primacor)
	carvedilol (Coreg)
	cardiac glycosides
	digitoxin
	digoxin
40	diuretics
	ACE Inhibitors
	Dobutamine
	dopamine
	<b>RESPIRATORY SYSTEM DISORDERS</b>
45	Asthma
	Chronic Obstructive Lung Disease (COLD) / Chronic Obstructive Pulmonary Disease (COPD)
	Acute Respiratory Distress Syndrome (ARDS)
50	Drug-Induced Pulmonary Disease
	Cystic Fibrosis

	Corticosteroids
	beclomethasone
	betamethasone
	cortisone
5	dexamethasone
	fluticasone (Flovent / Flonase)
	hydrocortisone
	methylprednisolone
	prednisolone
10	prednisone
	triamcinolone
	sympathomimetics
	albuterol (Proventil / Ventolin)
	salmeterol (Serevent)
15	muscarinic antagonists
	ipratropium (Combivent)
	leukotriene pathway inhibitors
	montelukast (Singulair)
	zafirtukast (Accolate)
20	mast cell stabilizers
	cromolyn (Intal)
	methylxanthines
	theophylline
	aminophylline
25	Dnase (Pulmozyme)

## **GASTROINTESTINAL SYSTEM DISORDERS**

	Gastro-esophageal Reflux Disease (GERD)
	Peptic Ulcer Disease
30	Inflammatory Bowel Disease
	Nausea and Vomiting
	Diarrhea, Constipation, Irritable Bowel Disease (IBD)
	Portal Hypertension and Cirrhosis
	Drug-Induced Liver Disease
35	Pancreatitis
	Viral Hepatitis
	Liver Transplantation
	Histamine-2 receptor antagonists
40	famotidine (Pepcid)
	nizatidine (Axid)
	pantoprazole (Protonix)
	rabeprazole (Aciphex)
	ranitidine (Zantac)
45	Proton Pump Inhibitors (PPIs)
	esomeprazole (Nexium)
	lansoprazole (Prevacid)
	omeprazole (Prilosec)
	Anti-nausea / anti-vertigo medicaments
50	anticholinergics

	antihistamines (Histamine-1 receptor antagonists)
	dopamine antagonists
	prokinetic gastric stimulant
	serotonin 5HT <sub>3</sub> receptor antagonists
5	dolasetron (Anzmet)
	granisetron (Kytril)
	ondansetron (Zofran)
	other medicaments
	hydroxyzine (Atarax, Vistaril)
10	corticosteroids
	benzodiazepines
	cannabinoids
	Prokinetic gastric stimulants (gastric motility stimulants)
	cisapride (Propulsid)
15	metoclopramide (Reglan)
	Laxatives
	Saline laxatives
	magnesium salts
	sodium salts
20	irritant / stimulant medicaments
	cascara
	senna
	phenolphthalein
	bisacodyl
25	casanthranol
	castor oil
	bulk-producing medicaments
	methylcellulose
	psyllium
30	polycarbophil
	lubricant
	mineral oil
	surfactants
	docusate
35	miscellaneous
	glycerin
	lactulose
	Anti-diarrheal medicaments
	diphenoxylate
40	atropine
	diphenoxin
	loperamide
	bismuth
	lactobacillus
45	Ulcerative Colitis Medicaments
	mesalamine
	olsalazine

## RENAL SYSTEM DISORDERS

50	Acute Renal Failure
----	---------------------

**NEUROLOGIC SYSTEM DISORDERS**

- 5 Multiple Sclerosis and inflammatory polyneuropathies  
Epilepsy  
Parkinson's disease and Movement Disorders  
Pain management  
Headache  
Amyotrophic Lateral Sclerosis
- 10 **Anti-epileptic medicaments**  
carbamazepine (Tegretol)  
divalproex sodium (Depakote)  
felbamate (Felbatol)  
15 gabapentin (Neurontin)  
lamotrigine (Lamictal)  
oxcarbazepine (Trileptal)  
phenytoin (Dilantin)  
topiramate (Topamax)  
20 zonisamide (Zonegran)
- Antimigraine medicaments**  
Serotonin 5HT<sub>1d</sub> receptor agonists  
almotriptan (Axert)  
frovatriptan (Frova)  
25 naratriptan (Amerge)  
rizatriptan (Rizalt)  
sumatriptan (Imitrex)  
zolmitriptan (Zomig)
- ergot alkaloids  
30 dihydroergotamine (DHE)  
isometheptine/dichlorophenazone (Midrin)  
caffeine  
pizotifen (Sanomigran)
- Sedative-hypnotic Medicaments**  
35 benzodiazepines  
alprazolam (Xanax)  
clonazepam (Kloriopin)  
clorazepate (Tranxene)  
diazepam (Valium)  
40 flumazenil (Romazicon) - antagonist  
lorazepam (Ativan)  
midazolam (Versed)  
triazolam (Halcion)
- barbiturates/Anesthetics  
45 pentobarbital (Nembutal)  
Phenobarbital (Luminal)  
thiopental (Pentothal)
- non-depressant anxiolytic  
buspirone (BuSpar)
- 50 **Treatment of alcoholism**

	disulfiram (Antabuse)
	<b>Pain Management Medicaments</b>
	Opioids
5	Opioid Peptides
	beta-endorphin
	dynorphin
	enkephalins
	Agonists
10	codeine
	etorphine
	fentanyl (Sublimaze)
	hydrocodeine
	hydromorphone
15	meperidine (Demerol)
	methadone (Dolophine)
	morphine
	oxycodone
	propoxyphene
20	Agonist-antagonists
	buprenorphine
	Partial Agonist
	dezocine (Dalgan)
	nalbuphine (Nubain)
	pentazocine (Talwain)
25	Antagonist
	naloxone (Narcan)
	Non-opiate
	acetaminophen (tylenol)
	tramadol (ultram)
30	<b>Anti-Parkinsonism Medicaments</b>
	levodopa
	carbidopa
	bromocriptine (Parlodel)
	pergolide (Permax)
35	amantadine (Symmetrel)
	selegiline (Deprenyl)
	anticholinergic agents
	dopamine Agonists
40	pramipexole (Mirapex)
	ropinirole (Requip)
	COMT inhibitors
	entacapone (Comtan)
	tolcapone (Tasmar)
45	<b>Anti-Spasticity Medicaments</b>
	baclofen (Lioresal)
	botulinum toxin type A (Botox)
	carisoprodol (Soma, Rela)
	chlorphenesin (Maolate)
	chlorzoxazone (Paraflex)
50	cyclobenzaprine (Flexeril)

- 5
  - dantrolene (Dantrium)
  - diazepam (Valium)
  - metaxalone (Skelaxin)
  - methocarbamol (Robaxin)
  - orphenadrine (Nor-flex)
  - tizanidine (Zanaflex)

## PSYCHIATRIC SYSTEM DISORDERS

- 10
  - Childhood psychiatric disorders
    - Attention Deficit Hyperactivity Disorder (ADHD) / Attention Deficit Disorder (ADD)
  - Eating disorders
  - Alzheimer's disease and Dementia Disorders
  - Substance abuse and Addictive Disorders
    - 15
      - alcohol, tobacco and caffeine abuse
  - Schizophrenia
  - Depressive disorders
  - Bipolar disorders
  - Anxiety disorders
  - 20
    - Obsessive-Compulsive disorders
    - Sleep disorders
- Psychostimulant Medicaments**
    - 25
      - amphetamine mixed salts (Adderall)
      - dextroamphetamine (Dexedrine)
      - methylphenidate (Ritalin, Concerta)
  - Antipsychotic Medicaments** (dopamine antagonists)
    - Phenothiazine type
      - 30
        - chlorpromazine (Thorazine)
        - fluphenazine (Prolixin)
    - Thioxanthene type
      - thiothixene (Navane)
    - Butyrophenone type
      - haloperidol (Haldol)
    - 35
      - Dibenzodiazepine type
        - clozapine (Clozaril)
      - Thienobenzodiazepine type
        - olanzapine (Zyprexa)
        - quetiapine (Seroquel)
    - 40
      - Antidepressant Medicaments**
        - Tricyclic antidepressants (TCA's)
          - amitriptyline (Elavil, Endep)
          - clomipramine (Anafranil), also a SSRI
          - desipramine (Norpramin)
          - 45
            - doxepin (Sinequan)
            - imipramine (Tofranil)
            - maprotiline (Ludiomil)
            - nortriptyline (Aventyl, Pamelor)
            - protriptyline (Vivactil)
        - 50
          - Monoamine oxidase inhibitors (MAO-I's)



	clorgyline (specific for MAO type A)
	isocarboxazid (Marplan)
	phenelzine (Nardil)
	tranylcypromine (Parnate)
5	Second Generation Medicaments (not including SSRIs)
	amoxapine (Asendin)
	bupropion (Wellbutrin)
	netazodone (Serzone)
	trazodone (Desyrel)
10	Serotonin-Specific Reuptake Inhibitors (SSRIs)
	citalopram (Celexa)
	clomipramine (Anafranil)
	escitalopram (Lexapro)
	fluoxetine (Prozac)
15	fluvoxamine (Luvox)
	paroxetine (Paxil)
	sertraline (Zoloft)
	Other
	lithium
20	mirtazapine (Temeron)
	venlafaxine (Effexor)
	<b>Anti-anxiety agents</b>
	barbiturates
	benzodiazepines
25	bupirone (Buspar) chloral hydrate
	doxepin
	hydroxyzine
	sedative-hypnotics
	serotonin reuptake inhibitors
30	<b>Anti-demential Medicaments</b>
	cholinesterase inhibitors
	donepezil (Aricept)
	galantamine (Reminyl)
	rivastigmine (Exelon)
35	tacrine (Cognex)

## ENDOCRINOLOGIC SYSTEM DISORDERS

	Diabetes mellitus
	Thyroid disorders
40	Adrenal Gland disorders
	Pituitary Gland disorders
	ACTH
	Adrenal androgens
45	Adrenocortical Function Antagonists
	Mineralocorticoid antagonists
	<b>Anti-Diabetic Medicaments</b>
	Insulin
50	Sulfonylureas

	acetoexamide (Dymelor)
	chlorpropamide (Diabinese)
	glimepiride (Amaryl)
5	glipizide (Glucotrol)
	glyburide (Micronase, DiaBeta)
	tolazamide (Tolinase)
	tolbutamide (Orinase)
	Biguanides
	metformin (Glucophage)
10	Alpha-glucosidase Inhibitors
	acarbose (Precose)
	miglitol (Glyset)
	Thiazolidinedione Derivatives
	pioglitazone (Actos)
15	rosiglitazone (Avandia)
	trogliatone (Rezulin)
	<b>Thyroid Disorder Medicaments</b>
	Levothyroxine
	Liothyronine
20	Liotrix
	<b>Hypothalamic and Pituitary Gland Medicaments</b>
	bromocriptine (Parlodel)
	chorionic gonadotropin (hCG)
	corticotropin generic (ACTH)
25	cosyntropin (Cortrosyn)
	desmopressin (DDAVP)
	gonadorelin acetate (GnRH) (Lutrepulse)
	gonadorelin hydrochloride (GnRH) (Factrel)
	goserelin acetate (Zoladex)
30	growth hormone
	histrelin (Supprelin)
	leuprolide (Lupron)
	menotropins (hMG) (Pergonal, Humegon)
	natarelin (Synarel)
35	octreotide (Sandostatin)
	oxytocin (Pitocinit, Syntocinon)
	pergolide (Permax)
	protirelin (Thybinone, Relefact TRH)
	sermorelin (GHRH) (Geref)
40	somatrem (Protropin)
	somatropin (Humatrope, Nutropin)
	thyrotropin (TSH) (Thytropar)
	urofollitropin (Metrodin)
	vasopressin (Pitressin Synthetic)

45	<b>GYNECOLOGIC SYSTEM AND OBSTETRIC CONDITIONS</b>
	Pregnancy and Lactation
	Infertility
	Contraception
50	Menstruation-related disorders

Endometriosis  
Hormone Replacement Therapy (HRT)

- 5           Conjugated estrogens (Premarin)  
            desogestrel  
            di-norgestrel  
            ethinyl diacetate  
            ethinyl estradiol  
            levonorgestrel  
10          medroxyprogesterone  
            norethindrone norgestimate  
            progesterone

#### **UROLOGIC SYSTEM DISORDERS**

- 15          Erectile Dysfunction  
            Benign Prostatic Hypertrophy  
            Urinary Incontinence
- 20          apomorphine  
            alprostadit  
            phosphodiesterase (PDE-5) inhibitors  
                sildenafil (Viagra)  
                tadalafil (Cialis)  
                vardenafil (Levitra)
- 25          tolterodine (Detrol)  
            tamulosin (Flomax)  
            yohimbine

#### **IMMUNOLOGIC SYSTEM DISORDERS**

- 30          Systemic Lupus Erythematosus and other Collagen-vascular diseases  
            Allergic and pseudo-allergic drug reactions

#### **BONE AND JOINT SYSTEM DISORDERS**

- 35          Osteoporosis and Osteomalacia  
            Rheumatoid Arthritis  
            Osteoarthritis  
            Gout and hyperuricemia

#### **Medicaments used in the Control of Inflammation**

- 40          Non-steroidal anti-inflammatory drugs (NSAIDs)  
            aspirin  
            diclofenac (Cataflam, Voltaren)  
            diflusnisal (Dolobid)  
            etodolac (Lodine)
- 45          fenoprofen (Nalfon)  
            flubiprofen (Ansaid)  
            ibuprofen (Motrin, Advil, Nuprin)  
            indomethacin (Indocin)  
            ketoprofen (Orudis)
- 50          ketorolac (Toradol)

	meclofenamate
	nabumetone (Relafen)
	naproxen (Naprosyn)
5	oxaprozin (Daypro)
	phenylbutazone
	piroxicam (Feldene)
	salicylate
	sulindac (Clinoril)
	tolmetin (Tolectin)
10	Cyclooxygenase-2 inhibitors (COX-2)
	celecoxib (Celebrex)
	rofecoxib (Vioxx)

#### **Arthritis and Gout Medicaments**

	allopurinol
15	anti-malarial compounds
	chloroquine
	colchicine
	enbrel
	Glucocorticoids
20	Gold
	methotrexate
	NSAIDs
	Penicillamine

#### **Other Medicaments**

25	alendronate (Fosamax)
	raloxifene (Evista)

### **DISORDERS OF THE EYES, EARS, NOSE, AND THROAT SYSTEMS**

30	Glaucoma
	Allergic rhinitis
	Histamine-1 receptor antagonists
	brompheniramine (Dimetane)
	cetirizine (Zyrtec)
35	chlorpheniramine (Chlor-Trimeton)
	clemastine (Tavist)
	cyproheptadine (Periactin)
	dimenhydrinate (Dramamine)
	diphenhydramine (Bendaryl)
40	doxylamine (Sominex, Unisom)
	fexofenadine (Allegra)
	loratidine (Claritin)
	Sympathomimetic medicaments
	pseudoephedrine (Sudated)
45	

### **DERMATOLOGIC SYSTEM DISORDERS**

	Acne
	Psoriasis
50	Rosacea and pigmentation disorders

## HEMATOLOGIC SYSTEM DISORDERS

- 5 Hematopoiesis
- Anemias
- Coagulation disorders
- Sickle-cell anemia
- Drug-induced hematologic disorders
- 10 **Coagulation disorders Medicaments**
  - aspirin
  - clopidogrel (Plavix)
  - fibrinolytic inhibitors
  - fibrinolytics
  - 15 glycoprotein (GP) IIb/IIIa antagonists / monoclonal antibodies
    - abciximab (Reopro)
    - eptifibatide (Integrelin)
    - tiofibran (Aggrastat)
  - heparin
  - 20 low-molecular weight heparins
  - Plasma fractions - blood factors
  - ticlopidine (Ticlid)
  - vitamin K
  - warfarin (Coumadin)

## INFECTIOUS SYSTEM DISEASES

- Central Nervous System (CNS) infections
- Lower Respiratory Tract Infections
- Upper Respiratory Tract Infections
- 30 Skin and Soft Tissue Infections
- Infective Endocarditis
- Tuberculosis
- Gastrointestinal Infections and Enterotoxigenic poisonings
- Intra-abdominal Infections
- 35 Parasitic diseases
- Urinary Tract Infections and Prostatitis
- Sexually Transmitted Diseases
- Bone and Joint Infections
- Sepsis and Septic Shock
- 40 Superficial Fungal Infections
- Invasive Fungal Infections
- Infections in Immunocompromised Patients
- Antimicrobial prophylaxis in Surgery
- Vaccines, toxoids, and other immunobiologics
- 45 Human Immunodeficiency Virus Infection

### Medicaments used in Infectious diseases

- Cell Wall Synthesis Inhibitors
  - Penicillins
  - 50 amoxicillin (Amoxil Polymox)

	ampicillin (Principen, Omnipen)
	benzathine Penicillin G
	benzyl Penicillin (Penicillin G)
5	carbenicillin (Geocillin)
	cloxacillin (Cloxapen)
	dicloxacillin (Dynapen)
	methicillin (Staphcillin)
	mezlocillin
10	nafcillin (Nafcil, Unipen)
	oxacillin
	phenoxymethyl Penicillin (Penicillin V)
	piperacillin (PiprachH)
	ticarcillin (Ticar)
	Cephalosporins
15	1st generation:
	cefazolin (Ancef, Defzol)
	cephalexin (Keflex)
	cephatothin (Keflin)
20	2nd generation:
	cefaclor (Ceclor)
	cefoxitin (Mefoxin)
	cefpodoxime (Vantin)
	cefuroxime (Zinacef, Ceftin)
	loracarbef (Lorabid)
25	3rd generation:
	cefoperazone
	cefotaxime (Claforan)
	cefotetan
30	ceftazidime (Fortax, Taxidime, Tazicef)
	ceftriaxone (Rocephin)
	veftizoxime (Cefizox)
	4th generation:
	cefepime
35	Other beta-Lactams aztreonam (Azactan)
	clavulanic acid
	imipenem (Primaxin)
	meropenem (Merrem IV)
	sulbactam
40	Other Cell-Wall Synthesis Inhibitors bacitracin
	cycloserine
	fosfomycin (Monurol)
	vancomycin (Vancocin)
	Agents Which Affect Cell Membranes
	Polymixins
45	Colistimethate
	Potymyxin B
	Protein Synthesis Inhibitors
	Aminoglycosides
50	amikacin (Amikin)
	gentamicin (Garamycin)

	kanamycin (Kantrex)
	neomycin
	netilmicin (Netromycin)
5	streptomycin
	tobramycin
	Tetracyclines
	demeclocycline (Declomycin)
	doxycycline
10	doxycycline (Vibramycin, Doryx)
	tetracycline (Achromycin)
	Macrolides
	azithromycin (Zithromax)
	clarithromycin (Biaxin)
15	erythromycin esters erythromycin
	Other Protein Synthesis Inhibitors
	Chloramphenicol (Chloromycetin)
	Clindamycin (Cleocin)
	Spectinomycin (Trobicin)
20	Inhibitors of Folate-Dependent Pathways
	co-trimoxazole
	silver Sulfadiazine
	sodium Sulfacetamide
	sulfamethoxazole (Gantanol)
25	sulfasalazine (Azulfidine) (Salicylazosulfapyridine)
	sulfisoxazole (Gantrisin)
	sulfonamides
	Dihydrofolate Reductase Inhibitor
	trimethoprim
30	DNA Gyrase Inhibitors
	ciprofloxacin (Cipro)
	gatifloxacin (Tequin)
	levofloxacin (Levaquin)
	lomefloxacin (Maxaquin)
35	nalidixic acid
	ofloxacin (Floxin)
	Urinary Tract Antiseptics
	nitrofurantoin
	Antimycobacterial Agents
40	First-line anti-TB medicaments
	ethambutol
	isoniazid (INI-I)
	pyrazinamide
	rifampin (Rimactane)
45	streptomycin
	Second-line anti-TB medicaments
	capreomycinA
	cycloserine
	dapsone
50	ethionamide
	para-aminosalicylic acid

	AntiFungal Agents
	amphotericin B (Fungizone, Amphotec)
	clotrimazole (Mycelex)
5	fluconazole (Diflucan)
	flucytosine
	griseofulvin
	itraconazole (Sporanox)
	ketoconazole (Nizoral)
10	miconazole (Monistat)
	nystatin (Mycostatin)
	AntiParasitic Agents
	Antimalarials
	chloroquine (Aralen)
15	mefloquine (Lariam)
	primaquine
	pyrimethamine-sulfadoxine (Fansidar)
	Anti protozoals
	metronidazole (Flagyl)
20	pentamidine isethionate
	pyrimethamine-sulfonamide
	trimethoprim (generic) sulfamethoxazole (Gantanol)
	Anthelmintic Medicaments
	mebendazole
25	praziquantel (Biltricide)
	pyrantel pamoate
	thiabendazole (Mintezol)
	Antiviral Medicaments
	acyclovir (Zovirax)
30	didanosine (DDI)
	foscarnet (Foscavir)
	ganciclovir (DHPG, Cytovene)
	ribavirin
	rimantadine
35	stavudine (d4T))
	valacyclovir (Valtrex)
	vidarabine (Vira-A)
	zalcitabine (ddC)
	zidovudine (Azidothymidine, AZT)
40	Protease inhibitors
	indinavir (Crixivan)
	ritonavir (Norvir)
	saquinavir (Fortovase)

## ONCOLOGIC AND IMMUNOLOGICAL DISORDERS

45	Breast Cancer
	Lung Cancer
	Colorectal Cancer
	Prostate Cancer
	Malignant Lymphomas
50	Ovarian Cancer



Acute Leukemias  
 Chronic Leukemias  
 Melanoma and other Skin Cancers  
 Hematopoeitic Stem Cell Transplantation

5

## **Anti-neoplastic Medicaments**

### **Alkylating Agents**

busulfan (Myleran)  
 carboplatin (Paraplatin)  
 10 carmustine (BNCU,BiCNU)  
 cisplatin (Platinol)  
 cyclophosphamide (Cytosan)  
 ifofamide (Ifex)  
 lomustine (CCNU,CeeNU)  
 15 mechlorethamine (Mustargen)  
 meiphalan (Alkeran)  
 procarbazine (Matulane)  
 thiotepa

### **Antimetabolites**

20 folic acid Antagonist  
 methotrexate  
 Purine Antagonists 6-mercaptopurine  
 6-thioguanine

### **Pyrimidine Antagonists**

25 cytarabine (ARA-C)  
 fluorouracil (5-FU)

### **Hormonal Agents: Hormones**

diethylstilbestrol (DES)  
 estrogens  
 30 prednisone (Deltasone)

### **Modulation of Hormone Release & Action Aminoglutethimide**

leuprolide acetate  
 tamoxifen (Nolvadex)

### **Plant Alkaloids**

35 Vinca Alkaloids  
 vinblastine (Velban)  
 vincristine (Oncovin)

### **Podophyllotoxins**

Etoposide (VP-16)

40 Other

docetaxel (Taxotere)  
 paclitaxel (Taxol)

### **Antibiotics**

45 bleomycin (Blenoxane)  
 dactinomycin (Cosmegen)  
 daunorubicin (DaunoXome)  
 doxorubicin (Adriamycin)  
 mitomycin (Mutamycin)

### **Other Anti-neoplastic Medicaments**

50 amsacrine (AMSA)

5	<ul style="list-style-type: none"> <li>azathioprine (Imuran)</li> <li>capecitabine (Xeloda)</li> <li>chlorambucil (Leukeran)</li> <li>cyclosporine (Sandimmune, Neoral)</li> <li>gemcitabine (Gemzar)</li> <li>hydroxyurea (Hydrea)</li> <li>mitotane (Sodren)</li> <li>mitoxantrone (Novantrone)</li> <li>pamidronate (Aredia)</li> </ul>
10	<b>Immunosuppressant Medicaments</b> <ul style="list-style-type: none"> <li>15-desoxyspergualin</li> <li>corticosteroids</li> <li>cyclosporine</li> <li>Interferons</li> </ul>
15	<ul style="list-style-type: none"> <li>Interleukins</li> <li>mycophenolate mofetil</li> <li>sirolimus (rapamycin)</li> <li>tacrolimus</li> <li>thalidomide</li> </ul>
20	<b>NUTRITIONAL DISORDERS</b> <ul style="list-style-type: none"> <li>Malnutrition, vitamin and mineral deficiencies</li> <li>Enteral Nutrition</li> <li>Obesity</li> </ul>
25	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>orlistat (Xenical)</li> <li>appetite suppressants</li> <li>sympathomimetic stimulants</li> <li>amphetamine stimulants</li> </ul> </li> <li>Mineral supplementation</li> </ul>
30	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>calcium ion</li> <li>iodine</li> <li>iron</li> <li>magnesium ion</li> <li>phosphorous</li> <li>potassium ion</li> </ul> </li> </ul>
35	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>selenium</li> <li>sodium ion</li> <li>zinc</li> </ul> </li> <li>Fat-soluble vitamins</li> </ul>
40	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>vitamin A</li> <li>vitamin D</li> <li>vitamin E</li> <li>vitamin K</li> </ul> </li> <li>Water-soluble vitamins</li> </ul>
45	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>vitamin C</li> <li>thiamine (vitamin B1)</li> <li>riboflavin (vitamin B2)</li> <li>niacin (vitamin B3)</li> <li>pyridoxine (vitamin B6)</li> <li>folate</li> </ul> </li> </ul>
50	<ul style="list-style-type: none"> <li> <ul style="list-style-type: none"> <li>cyanocobalamin (vitamin B12)</li> </ul> </li> </ul>

**MEDICAMENTS USED TO ALLEVIATE SYMPTOMS OF ALLERGIC  
RHINITIS, UPPER RESPIRATORY SYMPTOMS, COUGH, MILD ACHES  
AND PAINS**

5	Nasal Decongestants
	ephedrine
	phenylephrine
	phenylpropanolamine
	pseudoephedrine
10	Antihistamines (Histamine-1 receptor antagonists)
	Antitussive agents
	benzonatate
	codeine
	dextromethorphan
15	Expectorants
	guaifenesin
	iodinated glycerol
	terpin hydrate
	Xanthines
20	aminophylline
	caffeine
	dyphylline
	theophylline
	Pain relievers
25	narcotic agonists
	NSAIDS
	acetam inophen

**30      DIETARY SUPPLEMENTS**

	Arnica
	Bilberry
	Black Cohosh
	Cat's claw
35	Chamomile
	Echinacea
	Evening Primrose Oil
	Fenugreek
	Flaxseed
40	Feverfew
	Garlic
	Ginger root
	Ginkobiloba
	Ginseng
45	Goldenrod
	Hawthorn
	Kava-Kava
	Licorice
	Milk thistle
50	Psyllium

- 5
  - Rauwolfia
  - Senna
  - Soybean
  - St. John's wort
  - Saw palmetto
  - Turmeric
  - Valerian

## 10 **THERAPEUTIC PROTEINS and Biotechnology Medicaments**

**Additional Agents:** Norvasc, Neurontin, Paxil, Augmentin, Propecia, Lamisil, Lescol, bisphosphonate.

## 15 OTHER DRUGS

- abacavir sulfate
- acetazolamide
- acetylsalicylic acid
- 20 albendazole
- allopurinol
- amiloride hydrochloride
- amitriptyline hydrochloride
- artemether
- atropine sulfate
- 25 benznidazole
- biperiden hydrochloride
- chloroquine phosphate
- chlorpheniramine maleate
- chlorpromazine hydrochloride
- 30 cimetidine
- ciprofloxacin hydrochloride
- clofazimine
- clomiphene citrate
- clomipramine hydrochloride
- 35 cloxacillin sodium
- codeine phosphate
- dapsone
- didanosine
- diethylcarbamazine citrate
- 40 digoxin
- diloxanide furoate
- DL-methionine
- Doxycycline
- Efavirenz
- 45 ergometrine maleate
- ergotamine tartrate
- erythromycin ethyl succinate
- ethambutol hydrochloride
- ethosuximide
- 50 ferrous sulfate

	alendronate sodium
	amlodipine besylate
	amphetamine (mixed salts)
	atorvastatin calcium
5	benazepril hydrochloride
	bisoprolol fumarate
	bupropion hydrochloride
	carbidopa
	cefprozil
10	cetirizine hydrochloride
	citalopram hydrobromide
	clindamycin hydrochloride
	clonidine hydrochloride
	clopidogrel bisulfate
15	cyclobenzaprine hydrochloride
	desloratadine
	digoxin
	diltiazem hydrochloride
	doxazosin mesylate
20	doxycycline
	enalapril maleate
	fexofenadine hydrochloride
	fluoxetine hydrochloride
	folic acid
25	fosinopril sodium
	hydrocodone bitartrate
	hydrocodone
	hydroxyzine hydrochloride
	indinavir
30	irbesartan
	isosorbide mononitrate
	lamivudine
	levothyroxine sodium
	lopinavir
35	loratadine
	losartan potassium
	meclizine hydrochloride
	medroxyprogesterone acetate
	meperidine
40	metformin hydrochloride
	methylphenidate hydrochloride
	methylprednisolone
	metoclopramide hydrochloride)
	minocycline hydrochloride
45	montelukast sodium
	naproxen sodium
	nelfinavir
	nevirapine
	niclosamide
50	nicotinamide

	nifurtimox
	nitrofurantoin
	nortriptyline hydrochloride
	oxybutynin chloride
5	oxycodone hydrochloride
	paracetamol
	paroxetine hydrochloride
	penicillin V potassium
	phenytoin sodium
10	pioglitazone hydrochloride
	prednisolone
	primaquine phosphate
	pravastatin sodium
	prednisolone
15	promethazine hydrochloride
	promethazine fumarate
	propylthiouracil
	pyrantel embonate
	pyridostigmine bromide
20	raloxifene hydrochloride
	ranitidine hydrochloride
	rifampicin
	risedronate sodium
	risperidone
25	rosiglitazone maleate
	salbutamol sulfate
	saquinavir mesylate
	sertraline hydrochloride
	sildenafil citrate
30	sulfadiazine
	sumatriptan succinate
	tamoxifen citrate
	tamsulosin hydrochloride
	temazepam
35	terazosin hydrochloride
	timolol maleate
	tolterodine tartrate
	tramadol hydrochloride
	trazodone hydrochloride
40	triclabendazole
	valacyclovir hydrochloride
	valdecoxib
	valproic acid
	valsartan
45	venlafaxine hydrochloride
	verapamil hydrochloride
	warfarin sodium
	zolpidem tartrate

## EXAMPLES XII

As can be seen above, various embodiments of the present invention can be utilized in specific medical applications. By way of example only and not by way of limitation, the present invention can be practiced to prepare delivery devices for use in chemotherapy to address/treat, by way of example and not by limitation, the following aspects of chemotherapy: psychological, timing (to coincide with tumor growth for example) route of administration, nausea, vomiting (CINV), compliance, and cost (e.g. reduce hospital management of patients, reduce the number of “repeat” drug doses due to patient vomiting, etc.). Still further, the just mentioned aspects are not limited to chemotherapy, as the present invention can be practiced to address common aspects between chemotherapy and other treatments.

Further by way of examples, capsules containing Zofran (ondansertron), Temodar (temozolomide) can be made.

Still further, the present invention can be used in cardiovascular treatments, for example hypertension, heart failure, and heart rhythm disorders. Also, the present invention can be used in immunology (e.g. transplant rejections, auto-immune disorders, etc.). The present invention can be used to treat neurological disorders (such as Parkinson’s disease, dementia, stroke, epilepsy, and migraine headache, etc.), psychiatric disorders (schizophrenia, bipolar disease, depression, anxiety, ADHD / ADD, Addictions, etc.), infectious diseases (fungal, bacterial, viral (HIV), etc.), and in anesthesiology (induction anesthesia, local anesthesia). Furthermore, the present invention has application in endocrinology (cholesterol, diabetes, hormone replacement therapy, thyroid dysfunction, oral contraception, obesity, etc.), dermatology (onychomycosis, acne, rosacea, psoriasis, etc.), rheumatology (arthritis, gout, osteoporosis / Osteomalacia), respiratory fields (asthma, emphysema, cystic fibrosis, etc.), gastro-intestinal fields (gastro-esophageal reflux disease, ulcer prophylaxis, crohn’s disease, inflammatory bowel disease, etc.), chronic renal failure (vitamin and mineral replacement, blood pressure regulation, diabetes, depression, etc.), genito-urinary (enlarged prostate / BPH, overactive bladder, erectile dysfunction, feminine yeast infections, etc.) and

hematology-oncology (thromboembolous, hermatopoeisis, neoplastic disease, nausea / vomiting).

### EXAMPLES XIII

5

The present invention can be utilized with a variety of exipients. Categories of exipients include, but are not limited to, Binders, Disintegrants Fillers (diluents), Lubricants, Glidants (flow enhancers), Compression aids, Colors, Sweeteners, Preservatives, Suspending/dispersing agents, Film formers/coatings, Flavors, and Printing inks. Still further

10 by way of example and not by limitation, the present invention can be utilized with the following exipients:



Magnesium Stearate	Povidone
Lactose	Pregelatinized Starch
Microcrystalline Cellulose	Hydroxy Propyl Methylcellulose
Starch (corn)	OPA products (coatings & inks)
Silicon Dioxide	
Titanium Dioxide	Croscarmellose
Stearic Acid	Hydroxy Propyl Cellulose
Sodium Starch Glycolate	Ethylcellulose
Gelatin	Calcium Phosphate (dibasic)
Talc	Crospovidone
Sucrose	Shellac (and Glaze)
Calcium Stearate	

#### EXAMPLES XIV

- 5            Examples of the supporting nutraceutical formulations are to illustrate examples where specific categories of the natural products industry can be utilized with the present invention. There are many more categories than the ones that are listed and therefore this is for simply for the purpose to show that the technology is broad and could be utilized for many specific categories. The specific mg of each product is not included due to the amounts
- 10 of each material is typically based upon the formulators opinions, however there are some (RDA) recommended daily allowances that could be used to determine the formulation.

**Category:    Antioxidant**

- 15    Primary Capsule:

        d alpha Tocopherol  
        Beta Carotene  
        Tocotrineol

## Grape Seed Oil

### Secondary Capsule:

5           Selenium  
            Vitamin C Ester

### Category:    Brain Support

10

### Primary Capsule:

15           d alpha Tocopherol  
            DHA  
            Omega 3  
            Lecithin  
            Choline

### Secondary Capsule:

20

            Coenzyme Q 10  
            Ginkgo Biloba  
            B 12

### 25   Category:    Mood Support

### Primary Capsule:

30           D alpha Tocopherol  
            Lecithin  
            DHA  
            Omega 3

### Secondary Capsule:

35

            SAME  
            L Tyrosine

### 40   Category:    Cardio Support

### Primary Capsule:

45           d alpha Tocopherol  
            Tocotrienol  
            Flax Oil Omega 6  
            Fish Oil Omega 3

### Secondary Capsule:

50

Calcium  
Magnesium  
Coenzyme Q 10

5

**Category: Diet Support**

Primary Capsule:

10 Cojugated Linolic Acid  
Flax Seed Oil

Secondary Capsule:

15 Chromium  
Zinc  
L Carnitine

20 **Category: Immune Support**

Primary Capsule:

25 Garlic Oil  
Olive Leaf Oil  
d alpha Tocopherol

Secondary Capsule:

30 Zinc  
Echinacea

35 **Category: Laxative Support**

Primary Capsule:

40 Aloe Vera  
Flax Seed Oil

Secondary Capsule:

45 Senna Leaf  
Psyllium

**Category: Prostate Support**

50 Primary Capsule:

5 Saw Palmetto Oil  
Pygeum Oil  
Flaxseed Oil  
Pumpkin Seed Oil

Secondary Capsule:

10 Selenium  
Zinc  
Boswellia Serrata

15 **Category: Inflammation Support**

Primary Capsule:

20 Boswellia Serrata Oil  
Guggul Oil  
Omega 3 Oil  
Ginger Oil

Secondary Capsule:

25 Curcumin  
Holy Basil

30 **Category: Sports Nutrition / Muscle Support**

Primary Capsule:

35 Cojugated Linolic Acid  
MCT Oil

Secondary Capsule:

40 Zinc  
Chromium  
Tribulus Terrestris  
19 Nor-5-Androstendione

45 **Category: Menopause Support**

Primary Capsule:

50 Evening Primrose Oil

Red Raspberry Oil

Secondary Capsule:

5

Licorice Root  
Black Cohosh  
Soy Isoflavones

10 **Category: Cholesterol Support**

Primary Capsule:

15

Sterol Esters  
Guggul Oil  
d alpha Tocopherol  
Tocotrienol

20 Secondary Capsule:

Garlic Extract  
Zinc

25

\* \* \* \* \*

From the above discussion, it will be appreciated that the present invention provides novel integrated capsule delivery apparatus and methods for delivering diverse physical states (e.g., solid, liquid, gas or dispersion) of a single active ingredient or medicament (e.g., pharmaceutical, biotechnical, nutraceutical, vitamin, dietary supplement, mineral or combination thereof), or a plurality of active ingredients or medicaments, in a single dosage capsular form, wherein at least two of the active ingredients or medicaments if different receiving chambers have physical states that differ. In preferred design, the encapsulation processes and multi-compartment capsular technology of the present invention may include various desirable properties such as, for example, controlling time-release of key active ingredients or medicaments, prolonging shelf-life of the active ingredients or medicaments, improving palatability, reducing overall production costs and reducing the number of capsules consumed by a patient or consumer as nutritional or therapeutic agents.

The present invention provides novel integrated capsule delivery apparatus and methods for delivering a single dosage, multi-compartment capsule comprising a capsular base and cap configuration, wherein the size and shape of the cap, relative to its sealing relationship with the base, generally eliminates or substantially reduces any potential dead  
5 space volume within the internal periphery of the capsule, thereby functionally negating the opportunity for reaction between an air bubble and one or more active ingredients introduced into the capsule and, accordingly, improving stability of the capsular ingredient(s).

The present invention may be embodied in other specific forms without departing from its spirit or essential characteristics. The described embodiments are to be considered in  
10 all respects only as illustrative, and not restrictive. The scope of the invention is, therefore, indicated by the appended claims, rather than by the foregoing description. All changes which come within the meaning and range of equivalency of the claims are to be embraced within their scope.